

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 June 2003 (05.06.2003)


PCT

(10) International Publication Number
WO 03/045228 A2

- (51) International Patent Classification⁷: **A61B** (74) Agents: VINCENT, Matthew P. et al.; Ropes & Gray, One International Place, Boston, MA 02110-2624 (US).
- (21) International Application Number: PCT/US02/38347
- (22) International Filing Date:
26 November 2002 (26.11.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/333,691 26 November 2001 (26.11.2001) US
- (71) Applicants (for all designated States except US): TRUSTEES OF TUFTS COLLEGE [US/US]; Ballou Hall, Medford, MA 02155 (US). BRIGHAM AND WOMEN'S HOSPITAL [US/US]; 75 Francis Street, Boston, MA 02115 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): BACHOVCHIN, William W. [US/US]; 7 Warwick Road, Melrose, MA 02176 (US). KUCHROO, Vijay, K. [IN/US]; 30 Fairhaven Road, Newton, MA 02459-3424 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: METHODS FOR TREATING AUTOIMMUNE DISORDERS, AND REAGENTS RELATED THERETO



(57) Abstract: The invention generally relates to improved methods for treatment or prophylaxis in animal subjects (including humans) of autoimmune disorders including Type 1 diabetes, septic shock, multiple sclerosis, inflammatory bowel disease (IBD) and Crohn's disease.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

***Methods for Treating Autoimmune Disorders, and Reagents Related
Thereeto***

Funding

Work described herein was supported by funding from the National Institutes
5 of Health. The United States Government has certain rights in the invention.

Background of the Invention

The immune system can normally distinguish "self" from "non-self". Some
immune system cells (lymphocytes) become sensitized against "self" tissue cells, but
10 are normally controlled by other lymphocytes. When the normal control process is
disrupted, allowing lymphocytes to avoid suppression, or when there is an alteration
in some body tissue so that it is no longer recognized by the immune system as "self",
autoimmune disorders develop. The mechanisms that cause disrupted control or tissue
alterations are not well known. One theory holds that various microorganisms and
15 drugs may trigger some of these changes, particularly in people with genetic
predisposition to autoimmune disorders. There are a number of autoimmune diseases
including, for example, multiple sclerosis (MS), rheumatoid arthritis (RA), and Type
I diabetes.

Type I diabetes is a progressive autoimmune disease, in which the beta cells
20 that produce insulin are slowly destroyed by the body's own immune system. White
blood cells called T lymphocytes produce immune factors called cytokines that attack
and gradually destroy the beta cells of the pancreas. Important cytokines are
interleukin-1-beta, tumor necrosis factor-alpha, and interferon-gamma. Specific
proteins are also critical in the process. They include glutamic acid decarboxylase
25 (GAD), insulin, and islet cell antigens. These proteins serve as autoantigens. That is,
they trigger the self attack of the immune system on its body's own beta cells. It is
unknown what first starts this cascade of immune events, but evidence suggests that

both a genetic predisposition and environmental factors, such as a viral infection, are involved.

As a result of autoimmune diabetes, the pancreas produces little or no insulin, and insulin must be injected daily for the survival of the diabetic. Insulin, a hormone
5 produced by the pancreas, is needed to convert sugar, starches and other food into glucose and to make it available to the body's cells for energy. In muscle, adipose (fat) and connective tissues, insulin facilitates the entry of glucose into the cells by an action on the cell membranes. The ingested glucose is normally converted in the liver to CO₂ and H₂O (50%); to glycogen (5%); and to fat (30-40%), the latter being stored
10 in fat depots. Fatty acids from the adipose tissues are circulated, returned to the liver for re-synthesis of triacylglycerol and metabolized to ketone bodies for utilization by the tissues. The fatty acids are also metabolized by other organs. Fat formation is a major pathway for carbohydrate utilization. The net effect of insulin is to promote the storage and use of carbohydrates, protein and fat.

Some complications arising from long-standing diabetes are vascular disease, microvascular disease, eye complications, diabetic nephropathy, diabetic neuropathy, diabetic foot problems, and skin and mucous membrane problems. The action of Type 1 diabetes is to cause hyperglycemia (elevated blood glucose concentration) and a tendency towards diabetic ketoacidosis (DKA). Currently treatment requires chronic
15 administration of insulin. Sporadic or persistent incidence of hyperglycemia can be controlled by administering insulin. Uncontrolled hyperglycemia can further damage the cells of the pancreas which produce insulin (the β -islet cells) and in the long term create greater insulin deficiencies.
20

Type 1 diabetes (Insulin dependent diabetes mellitus, IDDM) represents 20%
25 of all human diabetes, and is the most serious form of the disease, with highest morbidity and mortality. Up to 800,000 people in the US are estimated to have type 1 diabetes, with about 30,000 new cases diagnosed each year. In addition, the incidence of IDDM has been rising over the past few decades in certain regions of the US and some European countries, particularly in Finland and England.

Currently, oral sulfonylureas and insulin injections are the only two therapeutic agents available in the United States for treatment of Diabetes mellitus. Both agents have the potential for producing hypoglycemia as a side effect, reducing the blood glucose concentration to dangerous levels. There is no generally applicable
5 and consistently effective means of maintaining an essentially normal fluctuation in glucose levels in DM. The resultant treatment attempts to minimize the risks of hypoglycemia while keeping the glucose levels below a target value. The drug regimen is combined with control of dietary intake of carbohydrates to keep glucose levels in control. However, to date, there has been no cure for many autoimmune
10 disorders, including type 1 diabetes. Clearly, a strong need exists for new, more effective treatments for these diseases.

Summary of the Invention

The invention generally relates to improved methods for treatment or
15 prophylaxis in animal subjects (including humans) of autoimmune disorders including Type I diabetes, septic shock, multiple sclerosis, inflammatory bowel disease (IBD) and Crohn's disease.

One aspect of the invention relates to a method for treating a mammal having an autoimmune disease, such as Type I diabetes, septic shock, multiple sclerosis, IBD,
20 or Crohn's disease, comprising administering to the mammal a dipeptidylpeptidase IV (DPIV) inhibitor, wherein administering the inhibitor induces immunosuppression and modulates the pharmacokinetics of polypeptide hormones required for tissue regeneration.

Brief Description of the Drawings

25

Figure 1 is a diagrammatic representation of the synthesis of a boro-Pro compound.

Figure 2 is a diagrammatic representation of the synthesis of a Cyclohexylglycine -boro-Ala.

Figure 3 is a time course of inactivation curve of Cyclohexylglycine-boro-Ala at pH 8.

5 Figure 4 is a bar graph illustrating DPPIV enzyme activity as measured from rat serum samples before and 1 hour after administration of Cyclohexylglycine-boro-Ala.

Figure 5 is a time course of incidence of Type 1 diabetes in NOD mice upon administration of Val-boro-Pro, or Cyclohexylglycine-boro-Ala, as compared to
10 control.

Figure 6 is a bar graph illustrating the incidence of Type 1 diabetes in NOD mice upon administration of Val-boro-Pro or Cyclohexylglycine-boro-Ala, as compared to control at 120 days after start of treatment.

15

Detailed Description of the Invention

i. Overview of the Invention

The present invention provides methods and compositions generally to prevent, reduce, or eliminate autoimmune disorders, such as Type 1 diabetes, septic
20 shock, multiple sclerosis, IBD or Crohn's disease in animals. As described in greater detail below, the subject method includes the administration, to an animal, of a composition including one or more dipeptidylpeptidase inhibitors, especially inhibitors of the dipeptidylpeptidase IV (DPIV) enzyme or other enzyme of similar specificity, in an amount which modulates the pharmacokinetics of polypeptide
25 hormones in a manner that produces tissue regeneration (e.g., such as may be manifest in an increase in tissue mass). While not being bound by any particular theory, it may be useful to deliver an amount of DPIV inhibitor sufficient to suppress an autoimmune component of the disease (such as a reduce a cellular and/or humoral

response against the target tissue). Other immunosuppressive agents can be administered conjointly.

For instance, in certain embodiments the method involves administration of a DPIV inhibitor, preferably at a predetermined time(s) during a 24-hour period, in an amount effective to improve one or more aberrant indices associated with Type 1 diabetes, septic shock, multiple sclerosis, IBD or Crohn's disease.

The present invention provides methods and compositions for regeneration of tissues damaged or destroyed as a result of autoimmune diseases by altering the half-life of a variety of different polypeptide hormones through inhibiting the proteolysis of one or more peptide hormones by DPIV or some other proteolytic activity.

The subject method can be used to increase the half-life of proglucagon-derived peptides including glucagon, glucagon-like peptide-1 (GLP-1), glucagon-like peptide-2 (GLP-2, PG 126-158), glicentin (corresponding to PG 1-69), oxyntomodulin (PG 33-69), glicentin-related pancreatic polypeptide (GRPP, PG 1-30), intervening peptide-2 (IP-2, PG 111-122amide), glucose-dependent insulintropic peptides, vasoactive intestinal polypeptide (VIP), vasostatin I and II, peptide histidine methionine (PHM), peptide histidine isoleucine (PHI), secretin, gastric inhibitory peptide, gastrin-releasing peptide (GRP), growth hormone-releasing hormone (GHRH), helospectin, helodermin, pituitary adenylate cyclase-activating peptide (PACAP, PACAP 27, and PACAP 38), and PACAP-related peptide (PRP), as well as calcitonin and secretin.

For instance, autoimmune (Type 1) diabetes leads to apoptotic destruction of β -pancreatic islets responsible for insulin secretion. Polypeptide hormones, such as exendin-4, an analog of glucagon-like peptide 1 (GLP-1), have been shown to stimulate the regeneration of the pancreas and expansion of β -cells mass by processes of both neogenesis and proliferation of β -cells. Support for β -cell neogenesis being a direct effect of the exendin-4 comes from the finding that duct cells expressed GLP-1 receptor, both at the mRNA and protein levels. (Xu et. al. (1999) Diabetes 48:2270-2276). Thus, modulation of plasma half-life of GLP-1 and analogs thereof hold

promise in a novel therapy to stimulate β -cell growth, and differentiation when administered to diabetic individuals with reduced β -cell mass.

5 In addition, GLP-1 and the longer acting GLP-1 agonist exendin-4 stimulate the expression of the IDX-1 homeodomain protein in the pancreas when administered to mice. Expression of both immunoreactive IDX-1 and β -galactosidase expressed from *LacZ* reporter transgene driven by the IDX-1 promoter occurs in the pancreatic ducts and the exocrine pancreas. It is also known that IDX-1 is required for the growth of the pancreas and that the epithelium of the pancreatic ducts and the centrolobular terminal ducts in the acinar tissue are the source of the differentiation of
10 new β -cells (β -cell neogenesis). (Stoffers et al (2000) Diabetes 49:741-748). Thus, modulation of plasma half-life of GLP-1 and analogs thereof similarly hold promise in a novel therapy to stimulate β -cell neogenesis.

Furthermore, the activation of inducible nitric oxide synthase (iNOS) by inflammatory cytokines is considered a mediator of destruction in β -cells. Recent
15 findings showed that the neuropeptide pituitary adenylyl cyclase-activating polypeptide (PACAP), whose distribution was identified in pancreatic neurons, inhibited nitric oxide (NO) production in cytokine-activated macrophages. (Sekiya (2000) Biochem Biophys Res Comm (2000) 278(1):211-216) Therefore, modulation of half-life of PACAP and analogs thereof may be useful in treatment of Type 1 diabetes.

20 GLP-2, for example, has been identified as a factor responsible for inducing proliferation of intestinal epithelium. See, for example, Drucker et al. (1996) PNAS 93:7911. The subject method can be used as part of a regimen for treating injury, inflammation, or resection of intestinal tissue, e.g., where enhanced growth and repair of the intestinal mucosal epithelial is desired. Thus modulation of half-life of GPL-2
25 and analogs thereof hold promise for the treatment of gut-related autoimmune diseases, such as, for example, inflammatory bowel disease (IBD).

Rheumatoid arthritis (RA) is a chronic and debilitating autoimmune disease of unknown etiology, characterized by chronic inflammation in the joints and subsequent destruction of the cartilage and bone. Neuropeptide vasoactive intestinal

peptide (VIP) has been shown to significantly reduce incidence and severity of arthritis, completely abrogating joint swelling and destruction of cartilage and bone. (Delgado (2001) Nat Med 7(5) 563-568) Consequently modulation of half-life of VIP and analogs thereof holds promise for the treatment of rheumatoid arthritis. In addition, VIP itself as well as more stable VIP derived agents, have been used or proposed as efficient therapeutic treatments of several autoimmune disorders, such as septic shock, multiple sclerosis, Crohn's disease, and autoimmune diabetes. (Gomariz (2001) Curr Pharm Des 7(2):89-111)

Additionally, the subject method can be used to alter the pharmacokinetics of Peptide YY (PYY) and neuropeptide Y (NPY), both members of the pancreatic polypeptide family, as DPIV has been implicated in the processing of those peptides in a manner which alters receptor selectivity.

DPIV has also been implicated in the metabolism and inactivation of growth hormone-releasing factor (GHRF). GHRF is a member of the family of homologous peptides that includes glucagon, secretin, vasoactive intestinal peptide (VIP), peptide histidine isoleucine (PHI), pituitary adenylate cyclase activating peptide (PACAP), gastric inhibitory peptide (GIP) and helodermin. Kubiak et al. (1994) Peptide Res 7:153. GHRF is secreted by the hypothalamus, and stimulates the release of growth hormone (GH) from the anterior pituitary. Thus, the subject method can be used to improve clinical therapy for certain growth hormone deficient children, and in clinical therapy of adults to improve nutrition and to alter body composition (muscle vs. fat). The subject method can also be used in veterinary practice, for example, to develop higher yield milk production and higher yield, leaner livestock.

Additional other autoimmune diseases and conditions that may be treated with the methods of the present invention are: systemic lupus erythematosus (SLE), immunovasculitis, myasthenia gravis, acute immunological arthritis, Hashimoto's thyroiditis, Grave's disease, rheumatoid synovitis, hereditary angioedema, hyperacute allograft rejection, xenograft rejection, infectious disease and sepsis, endotoxemia, scleroderma, glomerulonephritis, HIV infection, multiple sclerosis, atrophic gastritis, pancreatitis, allergic encephalomyelitis, thyrotoxicosis, sympathetic ophthalmia, allergic

reactions, multiple organ failure, allergic reactions, bullous diseases, urticaria, cryoglobulinemia, renal cortical necrosis, transplant organ reperfusion, post-ischemic reperfusion conditions, lupus nephritis, Crohn's disease, dermatomyositis, proliferative nephritis, type II collagen-induced arthritis, Behcet's syndrome, 5 Sjogren's syndrome, thermal injury, preeclampsia, thyroiditis, acute gouty arthritis, primary biliary cirrhosis inflammation, cranial nerve damage in meningitis, renal ischemia, anaphylaxis, and bowel inflammation.

In general, the inhibitors of the subject method will be small molecules, e.g., with molecular weights less than 7500 amu, preferably less than 5000 amu, and even 10 more preferably less than 2000 amu and even 1000 amu. In preferred embodiments, the inhibitors will be orally active.

In preferred embodiments, a DPIV inhibitor of the present invention has a K_i for DPIV inhibition of 1.0 nM or less, more preferably of 0.1 nM or less, and even more preferably of 0.01 nM or less. Indeed, inhibitors with K_i values in the picomolar 15 and even femtomolar range are contemplated.


Preferably, the compounds utilized in the subject method will have an EC_{50} for the desired biological effect, such as for example, increase in the plasma half-life of a peptide hormone that promotes tissue regeneration in the micromolar or less range more preferably in the nanomolar or less range and even more preferably in the 20 picomolar or femtomolar range.

In certain embodiments, the subject inhibitors are peptidyl compounds (including peptidomimetics) which are optimized, e.g., generally by selection of the $C\alpha$ substituents, for the substrate specificity of the targeted proteolytic activity. These peptidyl compounds will include a functional group, such as in place of the scissile 25 peptide bond, which facilitates inhibition of a serine-, cysteine- or aspartate-type protease, as appropriate. For example, the inhibitor can be a peptidyl α -diketone or a peptidyl α -keto ester, a peptide haloalkylketone, a peptide sulfonyl fluoride, a peptidyl boronate, a peptide epoxide, a peptidyl diazomethanes, a peptidyl phosphonate, isocoumarins, benzoxazin-4-ones, carbamates, isocyanates, isatoic 30 anhydrides or the like. Such functional groups have been provided in other protease

inhibitors, and general routes for their synthesis are known. See, for example, Angelastro et al., J. Med Chem. 33:11-13 (1990); Bey et al., EPO 363,284; Bey et al., EPO 364,344; Grubb et al., WO 88/10266; Higuchi et al., EPO 393,457; Ewoldt et al., Molecular Immunology 29(6):713-721 (1992); Hernandez et al., Journal of Medicinal Chemistry 35(6): 1121-1129 (1992); Vlasak et al., J Virology 63(5):2056-2062 (1989); Hudig et al., J Immunol 147(4):1360-1368 (1991); Odakc et al., Biochemistry 30(8):2217-2227 (1991); Vijayalakshmi et al., Biochemistry 30(8):2175-2183 (1991); Kam et al., Thrombosis and Haemostasis 64(1):133-137 (1990); Powers et al., J Cell Biochem 39(1):33-46 (1989); Powers et al., Proteinase Inhibitors, Barrett et al., Eds., Elsevier, pp. 55-152 (1986); Powers et al., Biochemistry 29(12):3108-3118 (1990); Oweida et al., Thrombosis Research 58(2):391-397 (1990); Hudig et al., Molecular Immunology 26(8):793-798 (1989); Orlowski et al., Archives of Biochemistry and Biophysics 269(1):125-136 (1989); Zunino et al., Biochimica et Biophysica Acta 967(3):331-340 (1988); Kam et al., Biochemistry 27(7):2547-2557 (1988); Parkes et al., Biochem J. 230:509-516 (1985); Green et al., J. Biol. Chem. 256:1923-1928 (1981); Angliker et al., Biochem. J. 241:871-875 (1987); Puri et al., Arch. Biochem. Biophys. 27:346-358 (1989); Hanada et al., Proteinase Inhibitors: Medical and Biological Aspects, Katunuma et al., Eds., Springer-Verlag pp. 25-36 (1983); Kajiwarra et al., Biochem. Int. 15:935-944 (1987); Rao et al., Thromb. Res. 47:635-637 (1987); Tsujinaka et al., Biochem. Biophys. Res. Commun. 153:1201-1208 (1988)). See also U.S. Patents Bachovchin et al. 4,935,493; Bachovchin et al. 5,462,928; Powers et al. 5,543,396; Hanko et al. 5,296,604; and the PCT publication of Ferring PCT/GB94/02615.

In other embodiments, the inhibitor is a non-peptidyl compound, e.g., which can be identified by such drug screening assays as described herein. These inhibitors can be, merely to illustrate, synthetic organic compounds, natural products, nucleic acids or carbohydrates.

A representative class of compounds for use in the method of the present invention are represented by the general formula (as described in the PCT application No. PCT/US99/02294, which is incorporated herein by reference):




wherein


A represents a 4-8-membered heterocycle including the N and the C α carbon;

5 Z represents C or N;

W represents a functional group which reacts with an active site residue of the targeted protease, as for example, -CN, -CH=NR₅,



R₁ represents a C-terminally linked amino acid residue or amino acid analog,
 10 or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or




R₂ is absent or represents one or more substitutions to the ring A, each of which can independently be a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido,
 15 -(CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-lower alkyl, -(CH₂)_m-O-lower alkenyl, -(CH₂)_n-O-(CH₂)_m-R₇, -(CH₂)_m-SH, -(CH₂)_m-S-lower alkyl, -(CH₂)_m-S-lower alkenyl, -(CH₂)_n-S-(CH₂)_m-R₇;

if X is N, R₃ represents hydrogen, if X is C, R₃ represents hydrogen or a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, -(CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-lower alkyl, -(CH₂)_m-O-lower alkenyl, -(CH₂)_n-O-(CH₂)_m-R₇, -(CH₂)_m-SH, -(CH₂)_m-S-lower alkyl, -(CH₂)_m-S-lower alkenyl, -(CH₂)_n-S-(CH₂)_m-R₇;

R₅ represents H, an alkyl, an alkenyl, an alkynyl, -C(X₁)(X₂)X₃, -(CH₂)_m-R₇, -(CH₂)_n-OH, -(CH₂)_n-O-alkyl, -(CH₂)_n-O-alkenyl, -(CH₂)_n-O-alkynyl, -(CH₂)_n-O-(CH₂)_m-R₇, -(CH₂)_n-SH, -(CH₂)_n-S-alkyl, -(CH₂)_n-S-alkenyl, -(CH₂)_n-S-alkynyl, -(CH₂)_n-S-(CH₂)_m-R₇, -C(O)C(O)NH₂, -C(O)C(O)OR'₇;

R₆ represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, -(CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-alkyl, -(CH₂)_m-O-alkenyl, -(CH₂)_m-O-alkynyl, -(CH₂)_m-O-(CH₂)_m-R₇, -(CH₂)_m-SH, -(CH₂)_m-S-alkyl, -(CH₂)_m-S-alkenyl, -(CH₂)_m-S-alkynyl, -(CH₂)_m-S-(CH₂)_m-R₇;



R₇ represents, independently for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R'₇ represents, independently for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

Y_1 and Y_2 can, independently, be OH, or a group capable of being hydrolyzed to a hydroxyl group, including cyclic derivatives where Y_1 and Y_2 taken together form a ring having from 5 to 8 atoms in the ring structure (such as pinacol or the like);

5 R_{50} represents O or S;

R_{51} represents N_3 , SH_2 , NH_2 , NO_2 or OR'_7 ;

R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which they are attached complete a heterocyclic ring having
10 from 5 to 8 atoms in the ring structure


X_1 represents a halogen;

X_2 and X_3 each independently represent a hydrogen or a halogen;

m is zero or an integer in the range of 1 to 8; and

n is an integer in the range of 1 to 8.

15 In preferred embodiments, the ring A is a 5, 6 or 7 membered ring, e.g., represented by the formula




and more preferably a 5- or 6-membered ring. The ring may, optionally, be further substituted.

20

In preferred embodiments, W represents $\begin{array}{c} Y_1 \\ | \\ -B- \\ | \\ Y_2 \end{array}$ or $\begin{array}{c} O \\ || \\ -C- \\ | \\ R_6 \end{array}$.

In preferred embodiments, R_1 is



wherein R_{36} is a small hydrophobic group, e.g., a lower alkyl or a halogen and R_{38} is hydrogen, or, R_{36} and R_{37} together form a 4-7-membered heterocycle including the N and the $C\alpha$ carbon, as defined for A above; and R_{40} represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group.

In preferred embodiments, R_2 is absent, or represents a small hydrophobic group such as a lower alkyl or a halogen.

In preferred embodiments, R_3 is a hydrogen, or a small hydrophobic group such as a lower alkyl or a halogen.


In preferred embodiments, R_5 is a hydrogen, or a halogenated lower alkyl.

In preferred embodiments, X_1 is a fluorine, and X_2 and X_3 , if halogens, are fluorine.

Also deemed as equivalents are any compounds which can be hydrolytically converted into any of the aforementioned compounds including boronic acid esters and halides, and carbonyl equivalents including acetals, hemiacetals, ketals, and hemiketals, and cyclic dipeptide analogs.


Longer peptide sequences are needed for the inhibition of certain proteases and improve the specificity of the inhibition in some cases.

In preferred embodiments, the subject method utilizes, as a DPIV inhibitor, a boronic acid analogs of an amino acid. For example, the present invention contemplates the use of boro-prolyl derivatives in the subject method. Exemplary boronic acid derived inhibitors of the present invention are represented by the general formula:




wherein

R_1 represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or



5

R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O\text{-alkyl}$, $-(CH_2)_m-O\text{-alkenyl}$, $-(CH_2)_m-O\text{-alkynyl}$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S\text{-alkyl}$, $-(CH_2)_m-S\text{-alkenyl}$, $-(CH_2)_m-S\text{-alkynyl}$, $-(CH_2)_m-S-(CH_2)_m-R_7$,



10

R_7 represents an aryl, a cycloalkyl, a cycloalkenyl, or a heterocycle;

R_8 and R_9 each independently represent hydrogen, alkyl, alkenyl, $-(CH_2)_m-R_7$, $-C(=O)\text{-alkyl}$, $-C(=O)\text{-alkenyl}$, $-C(=O)\text{-alkynyl}$, $-C(=O)\text{---(CH}_2)_m\text{---R}_7$, or R_8 and R_9 taken together with the N atom to which they are attached complete a heterocyclic ring having from 4 to 8 atoms in the ring structure;


15

R_{11} and R_{12} each independently represent hydrogen, a alkyl, or a pharmaceutically acceptable salt, or R_{11} and R_{12} taken together with the O-B-O atoms to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure;

5 m is zero or an integer in the range of 1 to 8; and


n is an integer in the range of 1 to 8.

In other embodiments, the subject DPIV inhibitors include an aldehyde analogs of proline or prolyl derivatives. Exemplary aldehyde-derived inhibitors of the present invention are represented by the general formula:




wherein

R_1 represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or



R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-alkynyl$, or $-(CH_2)_m-S-(CH_2)_m-R_7$.




R₇ represents an aryl, a cycloalkyl, a cycloalkenyl, or a heterocycle;

- R₈ and R₉ each independently represent hydrogen, alkyl, alkenyl, $\text{---(CH}_2\text{)}_m\text{---R}_7$,
 5 ---C(=O)---alkyl , $\text{---C(=O)---alkenyl}$, $\text{---C(=O)---alkynyl}$, $\text{---C(=O)---(CH}_2\text{)}_m\text{---R}_7$, or R₈ and R₉
 taken together with the N atom to which they are attached complete a heterocyclic
 ring having from 4 to 8 atoms in the ring structure; and

m is zero or an integer in the range of 1 to 8; and


n is an integer in the range of 1 to 8.

- 10 In yet further embodiments, the subject DPIV inhibitors are halo-methyl
 ketone analogs of an amino acid. Exemplary inhibitors of this class include
 compounds represented by the general formula:




wherein

- 15 R₁ represents a C-terminally linked amino acid residue or amino acid analog,
 or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or



R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-alkynyl$, or $-(CH_2)_m-S-(CH_2)_m-R_7$,



5

R_7 represents an aryl, a cycloalkyl, a cycloalkenyl, or a heterocycle;

R_8 and R_9 each independently represent hydrogen, alkyl, alkenyl, $-(CH_2)_m-R_7$, $-C(=O)-alkyl$, $-C(=O)-alkenyl$, $-C(=O)-alkynyl$, $-C(=O)-(CH_2)_m-R_7$, or R_8 and R_9


10 taken together with the N atom to which they are attached complete a heterocyclic ring having from 4 to 8 atoms in the ring structure;

X_1 , X_2 and X_3 each represent a hydrogen or a halogen; and


m is zero or an integer in the range of 1 to 8; and

n is an integer in the range of 1 to 8.

15 In preferred embodiments, the DPIV inhibitor is a peptide or peptidomimetic including a prolyl group or analog thereof in the P1 specificity position, and a nonpolar amino acid in the P2 specificity position, e.g., a nonpolar amino acid such as alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan or methionine, or an analog thereof. For example, the DPIV inhibitor may include an Ala-Pro or Pro-
 20 Pro dipeptide sequence or equivalent thereof, and be represented in the general formulas:



In preferred embodiments, the ring A is a 5-, 6-, or 7-membered ring, e.g., represented by the formula




- 5 In preferred embodiments, R_{32} is a small hydrophobic group, e.g., a lower alkyl or a halogen.

In preferred embodiments, R_{30} represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group.

- 10 In preferred embodiments, R_2 is absent, or represents a small hydrophobic group such as a lower alkyl or a halogen.


In preferred embodiments, R_3 is a hydrogen, or a small hydrophobic group such as a lower alkyl or a halogen.

- 15 Another representative class of compounds for use in the subject method include peptide and peptidomimetics of (D)-Ala-(L)-Ala, e.g., preserving the diastereomeric orientation of the substituents. Such inhibitors include compounds represented by the general formula:




wherein

W represents a functional group which reacts with an active site residue of the targeted protease, as for example, -CN, -CH=NR₅,




R₁ represents a C-terminally linked amino acid residue or amino acid analog,
5 or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or



R₃ represents hydrogen or a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an
10 acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, -(CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-lower alkyl, -(CH₂)_m-O-lower alkenyl, -(CH₂)_n-O-(CH₂)_m-R₇, -(CH₂)_m-SH, -(CH₂)_m-S-lower alkyl, -(CH₂)_m-S-lower alkenyl, -(CH₂)_n-S-(CH₂)_m-R₇;

R₅ represents H, an alkyl, an alkenyl, an alkynyl, -C(X₁)(X₂)X₃, -(CH₂)_m-R₇,
15 -(CH₂)_n-OH, -(CH₂)_n-O-alkyl, -(CH₂)_n-O-alkenyl, -(CH₂)_n-O-alkynyl, -(CH₂)_n-O-(CH₂)_m-R₇, -(CH₂)_n-SH, -(CH₂)_n-S-alkyl, -(CH₂)_n-S-alkenyl, -(CH₂)_n-S-alkynyl, -(CH₂)_n-S-(CH₂)_m-R₇, -C(O)C(O)NH₂, -C(O)C(O)OR'₇;

R₆ represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, -
(CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-alkyl, -(CH₂)_m-O-alkenyl, -(CH₂)_m-O-alkynyl,
20 -(CH₂)_m-O-(CH₂)_m-R₇, -(CH₂)_m-SH, -(CH₂)_m-S-alkyl, -(CH₂)_m-S-alkenyl, -(CH₂)_m-S-alkynyl, -(CH₂)_m-S-(CH₂)_m-R₇.



R₇ represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

- 5 R'₇ represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R₆₁ and R₆₂, independently, represent small hydrophobic groups;

- Y₁ and Y₂, independently, are OH, or a group capable of being hydrolyzed to a hydroxyl group, including cyclic derivatives where Y₁ and Y₂ taken together form a
 10 ring having from 5 to 8 atoms in the ring structure (such as pinacol or the like),

R₅₀ represents O or S;

R₅₁ represents N₃, SH₂, NH₂, NO₂ or OR'₇;

- R₅₂ represents hydrogen, a lower alkyl, an amine, OR'₇, or a pharmaceutically acceptable salt, or R₅₁ and R₅₂ taken together with the
 15 phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure

X₁ represents a halogen;

X₂ and X₃ each represent a hydrogen or a halogen

m is zero or an integer in the range of 1 to 8; and

n is an integer in the range of 1 to 8.

In preferred embodiments, R_1 is



wherein R_{36} is a small hydrophobic group, e.g., a lower alkyl or a halogen and R_{38} is hydrogen, or, R_{36} and R_{37} together form a 4-7 membered heterocycle including the N and the $C\alpha$ carbon, as defined for A above; and R_{40} represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group.

In preferred embodiments, R_3 is a hydrogen, or a small hydrophobic group such as a lower alkyl or a halogen.


In preferred embodiments, R_5 is a hydrogen, or a halogenated lower alkyl.

In preferred embodiments, X_1 is a fluorine, and X_2 and X_3 , if halogens, are fluorine.

In preferred embodiments, R_{61} and R_{62} , independently, represent low alkyls, such as methyl, ethyl, propyl, isopropyl, tert-butyl or the like.

Also included are such peptidomimetics as olefins, phosphonates, aza-amino acid analogs and the like.

Another representative class of compounds for use in the method of the present invention are represented by the general formula (as described in U.S. provisional application entitled "Methods for Regulating Glucose Metabolism, and Reagents Related Thereto" filed on Nov. 26, 2001, which is incorporated herein by reference):




wherein,

- 5 R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl, preferably lower alkyl such as methyl, ethyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl, preferably a bulky hydrophobic group, such as cyclohexyl, *t*-butyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

- 10 R_3 represents hydrogen or an amino-protecting group, preferably hydrogen;

R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or




- 15 R_6 represents hydrogen, a halogen, a alkyl, an alkenyl, an alkynyl, an aryl, $-(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-OH}$, $-(\text{CH}_2)_m\text{-O-alkyl}$, $-(\text{CH}_2)_m\text{-O-alkenyl}$, $-(\text{CH}_2)_m\text{-O-alkynyl}$, $-(\text{CH}_2)_m\text{-O-(CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-SH}$, $-(\text{CH}_2)_m\text{-S-alkyl}$, $-(\text{CH}_2)_m\text{-S-alkenyl}$, $-(\text{CH}_2)_m\text{-S-alkynyl}$, $-(\text{CH}_2)_m\text{-S-(CH}_2)_m\text{-R}_7$;

- 20 R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;


R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

U is absent or represents $-C(=O)-$, $-C(=S)-$, $-P(=O)(OR_8)-$, $-S(O_2)-$, or $-S(O)-$, preferably $-C(=O)-$, $-C(=S)-$, or $-S(O_2)-$;

W represents a functional group which reacts with an active site residue of the targeted protease, as for example, $-CN$, $-CH=NR_{53}$,



preferably



Y_1 and Y_2 are, independently, OH, or a group capable of being hydrolyzed, e.g., under physiologic conditions to a hydroxyl group, such as alkoxy, aryloxy, etc., including cyclic derivatives where Y_1 and Y_2 are connected via a ring having from 5 to 8 atoms in the ring structure (such as pinacol or the like);

R_{50} represents O or S;

R_{51} represents N_3 , SH, NH_2 , NO_2 or OR'_7 ;

R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure;

R_{53} represents hydrogen, an alkyl, an alkenyl, an alkynyl, $-C(X_1)(X_2)-X_3$, $-(CH_2)_m-R_7$, $-(CH_2)_n-OH$, $-(CH_2)_n-O\text{-alkyl}$, $-(CH_2)_n-O\text{-alkenyl}$, $-(CH_2)_n-O\text{-alkynyl}$, $-(CH_2)_n-O-(CH_2)_m-R_7$, $-(CH_2)_n-SH$, $-(CH_2)_n-S\text{-alkyl}$, $-(CH_2)_n-S\text{-alkenyl}$, $-(CH_2)_n-S\text{-alkynyl}$, $-(CH_2)_n-S-(CH_2)_m-R_7$, $-C(O)C(O)NH_2$, $-C(O)C(O)OR'_7$;


X_1 represents a halogen, preferably a fluorine;

X_2 and X_3 each represent a hydrogen or a halogen, preferably a hydrogen or a fluorine;

m is zero or an integer in the range of 1 to 8;

and n is an integer in the range of 1 to 8.

- 5 In preferred embodiments, the subject method utilizes, as a DPIV inhibitor, a boronic acid analog of an amino acid. For example, the present invention contemplates the use of boro-alanine derivatives in the subject method. Exemplary boronic acid derived inhibitors of the present invention are represented by the general formula:



10


wherein

R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl, preferably lower alkyl such as methyl, ethyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

- 15 R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl, preferably a bulky hydrophobic group, such as cyclohexyl, *t*-butyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

R_3 represents hydrogen or an amino-protecting group, preferably hydrogen;

- 20 R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or



preferably hydrogen.


R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, -
 $(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -alkyl, $-(CH_2)_m-O$ -alkenyl, $-(CH_2)_m-O$ -alkynyl, -
 5 $(CH_2)_m-O$ - $(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -alkyl, $-(CH_2)_m-S$ -alkenyl, $-(CH_2)_m-S$ -
 alkynyl, $-(CH_2)_m-S$ - $(CH_2)_m-R_7$;

R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl,
 cycloalkyl, cycloalkenyl, or heterocycle;

R_{11} and R_{12} each independently represent hydrogen, an alkyl, or a
 10 pharmaceutically acceptable salt, or R_{11} and R_{12} taken together with the O-B-O atoms
 to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in
 the ring structure; and

m is zero or an integer in the range of 1 to 8.

15 In other embodiments, the subject DPTV inhibitors include aldehyde analogs
 of alanine or alanyl derivatives. Exemplary aldehyde-derived inhibitors of the present
 invention are represented by the general formula:




wherein

20 R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower
 alkynyl, preferably lower alkyl such as methyl, ethyl, etc., optionally substituted by
 one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl, preferably a bulky hydrophobic group, such as cyclohexyl, *t*-butyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

5 R_3 represents hydrogen or an amino-protecting group, preferably hydrogen;

R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or



preferably hydrogen.


10

R_6 represents hydrogen, a halogen, a alkyl, an alkenyl, an alkynyl, an aryl, $-(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-OH}$, $-(\text{CH}_2)_m\text{-O-alkyl}$, $-(\text{CH}_2)_m\text{-O-alkenyl}$, $-(\text{CH}_2)_m\text{-O-alkynyl}$, $-(\text{CH}_2)_m\text{-O-(CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-SH}$, $-(\text{CH}_2)_m\text{-S-alkyl}$, $-(\text{CH}_2)_m\text{-S-alkenyl}$, $-(\text{CH}_2)_m\text{-S-alkynyl}$, $-(\text{CH}_2)_m\text{-S-(CH}_2)_m\text{-R}_7$;

15 R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

m is zero or an integer in the range of 1 to 8.

In yet further embodiments, the subject DPIV inhibitors are halo-methyl ketone analogs of an amino acid. Exemplary inhibitors of this class include
20 compounds represented by the general formula:




wherein,

R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl, preferably lower alkyl such as methyl, ethyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

- 5 R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl, preferably a bulky hydrophobic group, such as cyclohexyl, *t*-butyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

R_3 represents hydrogen or an amino-protecting group, preferably hydrogen;

- 10 R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or



preferably hydrogen.

- 15 R_6 represents hydrogen, a halogen, a alkyl, an alkenyl, an alkynyl, an aryl, -
 $(\text{CH}_2)_m\text{-}R_7$, $(\text{CH}_2)_m\text{-OH}$, $(\text{CH}_2)_m\text{-O-alkyl}$, $(\text{CH}_2)_m\text{-O-alkenyl}$, $(\text{CH}_2)_m\text{-O-alkynyl}$, -
 $(\text{CH}_2)_m\text{-O-(CH}_2)_m\text{-}R_7$, $(\text{CH}_2)_m\text{-SH}$, $(\text{CH}_2)_m\text{-S-alkyl}$, $(\text{CH}_2)_m\text{-S-alkenyl}$, $(\text{CH}_2)_m\text{-S-alkynyl}$, $(\text{CH}_2)_m\text{-S-(CH}_2)_m\text{-}R_7$;


- R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl,
 20 cycloalkyl, cycloalkenyl, or heterocycle;

X_1 , X_2 and X_3 each represent a hydrogen or a halogen;

m is zero or an integer in the range of 1 to 8.

In preferred embodiments, the DPIV inhibitor is a peptide or peptidomimetic including a alaninyl group or analog thereof in the P1 specificity position, and a non-naturally occurring amino acid in the P2 specificity position, or an analog thereof. For example, the DPIV inhibitor may include an Cyclohexylglycine-Ala or *t*-butylglycine-Ala dipeptide sequence or equivalent thereof, and be represented in the

5 general formula:




R₁ represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl, preferably lower alkyl such as methyl, ethyl, etc., optionally substituted by

10 one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

R₂ represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl, preferably a bulky hydrophobic group, such as cyclohexyl, *t*-butyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

15 R₃ represents hydrogen or an amino-protecting group, preferably hydrogen;

R₄ represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or




preferably hydrogen.

20


R₆ represents hydrogen, a halogen, a alkyl, an alkenyl, an alkynyl, an aryl, - (CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-alkyl, -(CH₂)_m-O-alkenyl, -(CH₂)_m-O-alkynyl, -

$(\text{CH}_2)_m\text{-O-(CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-SH}$, $-(\text{CH}_2)_m\text{-S-alkyl}$, $-(\text{CH}_2)_m\text{-S-alkenyl}$, $-(\text{CH}_2)_m\text{-S-alkynyl}$, $-(\text{CH}_2)_m\text{-S-(CH}_2)_m\text{-R}_7$;

R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle; W represents a functional group which reacts
5 with an active site residue of the targeted protease, as for example, $-\text{CN}$, $-\text{CH=NR}_{53}$,



preferably



Y_1 and Y_2 are, independently, OH , or a group capable of being hydrolyzed,
10 e.g., under physiologic conditions to a hydroxyl group, such as alkoxy, aryloxy, etc., including cyclic derivatives where Y_1 and Y_2 are connected via a ring having from 5 to 8 atoms in the ring structure (such as pinacol or the like);

R_{50} represents O or S;

R_{51} represents N_3 , SH , NH_2 , NO_2 or OR'_7 ;

R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure;


R_{53} represents hydrogen, an alkyl, an alkenyl, an alkynyl, $-\text{C}(\text{X}_1)(\text{X}_2)-\text{X}_3$,
20 $(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_n\text{-OH}$, $-(\text{CH}_2)_n\text{-O-alkyl}$, $-(\text{CH}_2)_n\text{-O-alkenyl}$, $-(\text{CH}_2)_n\text{-O-alkynyl}$, $-(\text{CH}_2)_n\text{-O-(CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_n\text{-SH}$, $-(\text{CH}_2)_n\text{-S-alkyl}$, $-(\text{CH}_2)_n\text{-S-alkenyl}$, $-(\text{CH}_2)_n\text{-S-alkynyl}$, $-(\text{CH}_2)_n\text{-S-(CH}_2)_m\text{-R}_7$, $-\text{C}(\text{O})\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{C}(\text{O})\text{OR}'_7$, preferably a hydrogen, or a halogenated lower alkyl;

X_1 represents a halogen, preferably a fluorine;

m is zero or an integer in the range of 1 to 8; and

n is an integer in the range of 1 to 8.

- Another representative class of compounds for use in the subject method include peptide and peptidomimetics of (L)-Ala-(L)-Cyclohexylglycine, e.g., preserving the steric disposition of moieties. Such inhibitors include compounds represented by the general formula:




wherein,

- R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl, preferably lower alkyl such as methyl, ethyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

- R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl, preferably a bulky hydrophobic group, such as cyclohexyl, *t*-butyl, etc., optionally substituted by one or more small substituents such as halogen, hydroxy, alkoxy, etc.;

R_3 represents hydrogen or an amino-protecting group, preferably hydrogen;

- R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or




preferably hydrogen.


R_6 represents hydrogen, a halogen, a alkyl, an alkenyl, an alkynyl, an aryl, -
 $(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, -
 $(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-$
 5 alkynyl, $-(CH_2)_m-S-(CH_2)_m-R_7$;

R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl,
 cycloalkyl, cycloalkenyl, or heterocycle;

W represents a functional group which reacts with an active site residue of the
 targeted protease, as for example, $-CN$, $-CH=NR_{53}$,



preferably



Y_1 and Y_2 are, independently, OH , or a group capable of being hydrolyzed,
 e.g., under physiologic conditions to a hydroxyl group, such as alkoxy, aryloxy, etc.,
 15 including cyclic derivatives where Y_1 and Y_2 are connected via a ring having from 5
 to 8 atoms in the ring structure (such as pinacol or the like);

R_{50} represents O or S ;

R_{51} represents N_3 , SH , NH_2 , NO_2 or OR'_7 ;

R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically
 20 acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which
 they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring
 structure;

R_{53} represents hydrogen, an alkyl, an alkenyl, an alkynyl, $-C(X_1)(X_2)-X_3$, $-(CH_2)_m-R_7$, $-(CH_2)_n-OH$, $-(CH_2)_n-O-alkyl$, $-(CH_2)_n-O-alkenyl$, $-(CH_2)_n-O-alkynyl$, $-(CH_2)_n-O-(CH_2)_m-R_7$, $-(CH_2)_n-SH$, $-(CH_2)_n-S-alkyl$, $-(CH_2)_n-S-alkenyl$, $-(CH_2)_n-S-alkynyl$, $-(CH_2)_n-S-(CH_2)_m-R_7$, $-C(O)C(O)NH_2$, $-C(O)C(O)OR_7$, preferably a
5 hydrogen, or a halogenated lower alkyl;

X_1 represents a halogen, preferably a fluorine;

m is zero or an integer in the range of 1 to 8;

and n is an integer in the range of 1 to 8.

Also deemed as equivalents are any compounds which can be hydrolytically
10 converted into any of the aforementioned compounds including boronic acid esters and halides, and carbonyl equivalents including acetals, hemiacetals, ketals, and hemiketals, and cyclic dipeptide analogs.

As used herein, the definition of each expression, e.g. alkyl, m , n , etc., when it
occurs more than once in any structure, is intended to be independent of its definition
15 elsewhere in the same structure.

The pharmaceutically acceptable salts of the subject compounds include the
conventional nontoxic salts or quaternary ammonium salts of the compounds, e.g.,
from non-toxic organic or inorganic acids. For example, such conventional nontoxic
salts include those derived from inorganic acids such as hydrochloric, hydrobromic,
20 sulfuric, sulfamic, phosphoric, nitric, and the like; and the salts prepared from organic
acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric,
ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic,
sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane
disulfonic, oxalic, isothionic, and the like.

25 The pharmaceutically acceptable salts of the present invention can be
synthesized from the subject compound which contain a basic or acid moiety by
conventional chemical methods. Generally, the salts are prepared by reacting the free
base or acid with stoichiometric amounts or with an excess of the desired salt-forming
inorganic or organic acid or base in a suitable solvent. The pharmaceutically

acceptable salts of the acids of the subject compounds are also readily prepared by conventional procedures such as treating an acid with an appropriate amount of a base such as an alkali or alkaline earth methyl hydroxide (e.g. sodium, potassium, lithium, calcium or magnesium) or an organic base such as an amine, piperidine, pyrrolidine, benzylamine and the like, or a quaternary ammonium hydroxide such as tetramethylammonium hydroxide and the like.

Contemplated equivalents of the compounds described above include compounds which otherwise correspond thereto, and which have the same general properties thereof (e.g. the ability to inhibit proteolysis of GLP-1 or other peptide hormone or precursor thereof), wherein one or more simple variations of substituents are made which do not adversely affect the efficacy of the compound in use in the contemplated method. In general, the compounds of the present invention may be prepared by the methods illustrated in the general reaction schemes as, for example, described below, or by modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are in themselves known, but are not mentioned here.

In other embodiments of the methods, the present invention further contemplates the use of known DPIV inhibitors in the art, such as, for example, TMC-2A, TMC-2B and TMC-2C (Nonaka (1997) J. Antibiot (Tokyo) 50(8):646-652); Lys[Z(NO₂)]-thiazolidide, Lys[Z(NO₂)]-piperidide, and Lys[Z(NO₂)]-pyrrolidide (Reinhold et al. (1997) Immunology 91(3):354-360); Phenylalanyl-pyrrolidine-2-nitrile and arginyl(PMC)-pyrrolidine-2-nitrile (Jiang et al (1997) Res. Virol. 148(4):255-266); Ala-Pro-nitrobenzoylhydroxylamine (Tanaka et al (1997) Int J Immunopharmacol 19(1):15-24; Ala-PipP(OPh-4-Cl)₂, Ala-ProP(OPh)₂, Ala-ProP(OPh-4-Cl)₂, (Boduszek et al (1994) J Med Chem 37(23):3969-3976; diprotin A and diprotin B (Umezawa et al (1984) J Antibiotics 37:422-425); 4-amino-(2,6-dimethylphenyl)phthalimides, 4- and 5-hydroxy-(2,6-diethylphenyl)phthalimide, and 4-hydroxy-(2,6-diisopropylphenyl)phthalimide (Shimazawa et al (1999) Bioorg Med Chem Lett 9(4):559-562). Latest developments in the search of DPIV inhibitors have

also been reviewed (Augustyns et al (1999) Curr Med Chem 6(4):311-327). All of the above-cited references and publications are hereby incorporated by reference.

ii. Definitions

5 For convenience, before further description of the present invention, certain terms employed in the specification, examples, and appended claims are collected here.

The term "alkyl" refers to the radical of saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In preferred embodiments, a straight chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (e.g., C₁-C₃₀ for straight chain, C₃-C₃₀ for branched chain), and more preferably 20 or fewer. Likewise, preferred cycloalkyls have from 3-10 carbon atoms in their ring structure, and more preferably have 5, 6 or 7 carbons in the ring structure.

Moreover, the term "alkyl" (or "lower alkyl") as used throughout the specification and claims is intended to include both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, a halogen, a hydroxyl, a carbonyl (such as a carboxyl, an ester, a formyl, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an alkoxyl, a phosphoryl, a phosphonate, a phosphinate, an amino, an amido, an amidine, an imine, a cyano, a nitro, an azido, a sulfhydryl, an alkylthio, a sulfate, a sulfonate, a sulfamoyl, a sulfonamido, a sulfonyl, a heterocyclyl, an aralkyl, or an aromatic or heteroaromatic moiety. It will be understood by those skilled in the art that the moieties substituted on the hydrocarbon chain can themselves be substituted, if appropriate. For instance, the substituents of a substituted alkyl may include substituted and unsubstituted forms of amino, azido, imino, amido, phosphoryl (including phosphonate and phosphinate), sulfonyl (including sulfate,

sulfonamido, sulfamoyl and sulfonate), and silyl groups, as well as ethers, alkylthios, carbonyls (including ketones, aldehydes, carboxylates, and esters), -CF₃, -CN and the like. Exemplary substituted alkyls are described below. Cycloalkyls can be further substituted with alkyls, alkenyls, alkoxys, alkylthios, aminoalkyls, carbonyl-substituted alkyls, -CF₃, -CN, and the like.

The term "aralkyl", as used herein, refers to an alkyl group substituted with an aryl group (e.g., an aromatic or heteroaromatic group).

The terms "alkenyl" and "alkynyl" refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively.

Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to ten carbons, more preferably from one to six carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths. Preferred alkyl groups are lower alkyls. In preferred embodiments, a substituent designated herein as alkyl is a lower alkyl.

The term "aryl" as used herein includes 5-, 6- and 7-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, triazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles" or "heteroaromatics". The aromatic ring can be substituted at one or more ring positions with such substituents as described above, for example, halogen, azide, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, sulfonamido, ketone, aldehyde, ester, heterocyclyl, aromatic or heteroaromatic moieties, -CF₃, -CN, or the like. The term "aryl" also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings (the rings are "fused rings") wherein at least one of the rings is

aromatic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls.

The terms "heterocyclyl" or "heterocyclic group" refer to 3- to 10-membered ring structures, more preferably 3- to 7-membered rings, whose ring structures include one to four heteroatoms. Heterocycles can also be polycycles. Heterocyclyl groups include, for example, thiophene, thianthrene, furan, pyran, isobenzofuran, chromene, xanthene, phenoxathiin, pyrrole, imidazole, pyrazole, isothiazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, pyrimidine, phenanthroline, phenazine, phenarsazine, phenothiazine, furazan, phenoxazine, pyrrolidine, oxolane, thiolane, oxazole, piperidine, piperazine, morpholine, lactones, lactams such as azetidinones and pyrrolidinones, sultams, sultones, and the like. The heterocyclic ring can be substituted at one or more positions with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.


The terms "polycyclyl" or "polycyclic group" refer to two or more rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings, e.g., the rings are "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle can be substituted with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.

The term "carbocycle", as used herein, refers to an aromatic or non-aromatic ring in which each atom of the ring is carbon.

The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms are nitrogen, oxygen, sulfur and phosphorous.

As used herein, the term "nitro" means -NO_2 ; the term "halogen" designates -F, -Cl, -Br or -I; the term "sulfhydryl" means -SH ; the term "hydroxyl" means -OH ; and the term "sulfonyl" means $\text{-SO}_2\text{-}$.

The terms "amine" and "amino" are art recognized and refer to both unsubstituted and substituted amines, e.g., a moiety that can be represented by the general formula:




10

wherein R_9 , R_{10} and R'_{10} each independently represent a hydrogen, an alkyl, an alkenyl, $\text{-(CH}_2\text{)}_m\text{-R}_8$, or R_9 and R_{10} taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure; R_8 represents an aryl, a cycloalkyl, a cycloalkenyl, a heterocycle or a polycycle; and m is zero or an integer in the range of 1 to 8. In preferred embodiments, only one of R_9 or R_{10} can be a carbonyl, e.g., R_9 , R_{10} and the nitrogen together do not form an imide. In even more preferred embodiments, R_9 and R_{10} (and optionally R'_{10}) each independently represent a hydrogen, an alkyl, an alkenyl, or $\text{-(CH}_2\text{)}_m\text{-R}_8$. Thus, the term "alkylamine" as used herein means an amine group, as defined above, having a substituted or unsubstituted alkyl attached thereto, i.e., at least one of R_9 and R_{10} is an alkyl group.


20

The term "acylamino" is art-recognized and refers to a moiety that can be represented by the general formula:



wherein R₉ is as defined above, and R'₁₁ represents a hydrogen, an alkyl, an alkenyl or $\text{---}(\text{CH}_2)_m\text{---R}_8$, where m and R₈ are as defined above.


The term "amido" is art recognized as an amino-substituted carbonyl and includes a moiety that can be represented by the general formula:



wherein R₉, R₁₀ are as defined above. Preferred embodiments of the amide will not include imides which may be unstable.

The term "alkylthio" refers to an alkyl group, as defined above, having a sulfur radical attached thereto. In preferred embodiments, the "alkylthio" moiety is represented by one of ---S---alkyl , ---S---alkenyl , ---S---alkynyl , and $\text{---S---}(\text{CH}_2)_m\text{---R}_8$, wherein m and R₈ are defined above. Representative alkylthio groups include methylthio, ethylthio, and the like.

The term "carbonyl" is art recognized and includes such moieties as can be represented by the general formula:



wherein X is a bond or represents an oxygen or a sulfur, and R₁₁ represents a hydrogen, an alkyl, an alkenyl, $\text{---}(\text{CH}_2)_m\text{---R}_8$ or a pharmaceutically acceptable salt, R'₁₁ represents a hydrogen, an alkyl, an alkenyl or $\text{---}(\text{CH}_2)_m\text{---R}_8$, where m and R₈ are as defined above. Where X is an oxygen and R₁₁ or R'₁₁ is not hydrogen, the

formula represents an "ester". Where X is an oxygen, and R₁₁ is as defined above, the moiety is referred to herein as a carboxyl group, and particularly when R₁₁ is a hydrogen, the formula represents a "carboxylic acid". Where X is an oxygen, and R'₁₁ is hydrogen, the formula represents a "formate". In general, where the oxygen


5 atom of the above formula is replaced by sulfur, the formula represents a "thiolcarbonyl" group. Where X is a sulfur and R₁₁ or R'₁₁ is not hydrogen, the formula represents a "thioester." Where X is a sulfur and R₁₁ is hydrogen, the formula represents a "thiocarboxylic acid." Where X is a sulfur and R'₁₁ is hydrogen, the formula represents a "thioformate." On the other hand, where X is a

10 bond, and R₁₁ is not hydrogen, the above formula represents a "ketone" group. Where X is a bond, and R₁₁ is hydrogen, the above formula represents an "aldehyde" group.

The terms "alkoxyl" or "alkoxy" as used herein refers to an alkyl group, as defined above, having an oxygen radical attached thereto. Representative alkoxyl

15 groups include methoxy, ethoxy, propyloxy, tert-butoxy and the like. An "ether" is two hydrocarbons covalently linked by an oxygen. Accordingly, the substituent of an alkyl that renders that alkyl an ether is or resembles an alkoxyl, such as can be represented by one of -O-alkyl, -O-alkenyl, -O-alkynyl, -O-(CH₂)_m-R_g, where m and R_g are described above.


20 The term "sulfonate" is art recognized and includes a moiety that can be represented by the general formula:



in which R₄₁ is an electron pair, hydrogen, alkyl, cycloalkyl, or aryl.


The term "sulfate" is art recognized and includes a moiety that can be

25 represented by the general formula:



in which R_{41} is as defined above.


The term "sulfonamido" is art recognized and includes a moiety that can be represented by the general formula:



5


in which R_9 and R'_{11} are as defined above.

The term "sulfamoyl" is art-recognized and includes a moiety that can be represented by the general formula:




10 in which R_9 and R_{10} are as defined above.

The terms "sulfoxido" or "sulfinyl", as used herein, refers to a moiety that can be represented by the general formula:




in which R_{44} is selected from the group consisting of hydrogen, alkyl, alkenyl,
15 alkynyl, cycloalkyl, heterocyclyl, aralkyl, or aryl.

A "phosphoryl" can in general be represented by the formula:




- wherein Q_1 represented S or O, and R_{46} represents hydrogen, a lower alkyl or an aryl. When used to substitute, e.g., an alkyl, the phosphoryl group of the
- 5 phosphorylalkyl can be represented by the general formula:




wherein Q_1 represented S or O, and each R_{46} independently represents hydrogen, a lower alkyl or an aryl, Q_2 represents O, S or N. When Q_1 is an S, the phosphoryl moiety is a "phosphorothioate".

- 10 A "phosphoramidite" can be represented in the general formula:



wherein R_9 and R_{10} are as defined above, and Q_2 represents O, S or N.

A "phosphonamidite" can be represented in the general formula:



- 15 wherein R_9 and R_{10} are as defined above, Q_2 represents O, S or N, and R_{48} represents a lower alkyl or an aryl, Q_2 represents O, S or N.

A "selenoalkyl" refers to an alkyl group having a substituted seleno group attached thereto. Exemplary "selenoethers" which may be substituted on the alkyl are

selected from one of -Se-alkyl, -Se-alkenyl, -Se-alkynyl, and -Se-(CH₂)_m-R₇, m and R₇ being defined above.

Analogous substitutions can be made to alkenyl and alkynyl groups to produce, for example, aminoalkenyls, aminoalkynyls, amidoalkenyls, amidoalkynyls, iminoalkenyls, iminoalkynyls, thioalkenyls, thioalkynyls, carbonyl-substituted alkenyls or alkynyls.

It will be understood that "substitution" or "substituted with" includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc.

As used herein, the term "substituted" is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. Illustrative substituents include, for example, those described hereinabove. The permissible substituents can be one or more and the same or different for appropriate organic compounds. For purposes of this invention, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valencies of the heteroatoms. This invention is not intended to be limited in any manner by the permissible substituents of organic compounds.

A "small" substituent is one of 10 atoms or less.

By the terms "amino acid residue" and "peptide residue" is meant an amino acid or peptide molecule without the -OH of its carboxyl group. In general the abbreviations used herein for designating the amino acids and the protective groups are based on recommendations of the IUPAC-IUB Commission on Biochemical Nomenclature (see Biochemistry (1972) 11:1726-1732). For instance Met, Ile, Leu, Ala and Gly represent "residues" of methionine, isoleucine, leucine, alanine and

glycine, respectively. By the residue is meant a radical derived from the corresponding α -amino acid by eliminating the OH portion of the carboxyl group and the H portion of the α -amino group. The term "amino acid side chain" is that part of an amino acid exclusive of the $-\text{CH}(\text{NH}_2)\text{COOH}$ portion, as defined by K. D.

5 Kopple, "Peptides and Amino Acids", W. A. Benjamin Inc., New York and Amsterdam, 1966, pages 2 and 33; examples of such side chains of the common amino acids are $-\text{CH}_2\text{CH}_2\text{SCH}_3$ (the side chain of methionine), $-\text{CH}_2(\text{CH}_3)-\text{CH}_2\text{CH}_3$ (the side chain of isoleucine), $-\text{CH}_2\text{CH}(\text{CH}_3)_2$ (the side chain of leucine) or H-(the side chain of glycine).

10 For the most part, the amino acids used in the application of this invention are those naturally occurring amino acids found in proteins, or the naturally occurring anabolic or catabolic products of such amino acids which contain amino and carboxyl groups. Particularly suitable amino acid side chains include side chains selected from those of the following amino acids: glycine, alanine, valine, cysteine, leucine,
15 isoleucine, serine, threonine, methionine, glutamic acid, aspartic acid, glutamine, asparagine, lysine, arginine, proline, histidine, phenylalanine, tyrosine, and tryptophan, and those amino acids and amino acid analogs which have been identified as constituents of peptidylglycan bacterial cell walls.

The term amino acid residue further includes analogs, derivatives and
20 congeners of any specific amino acid referred to herein, as well as C-terminal or N-terminal protected amino acid derivatives (e.g. modified with an N-terminal or C-terminal protecting group). For example, the present invention contemplates the use of amino acid analogs wherein a side chain is lengthened or shortened while still providing a carboxyl, amino or other reactive precursor functional group for
25 cyclization, as well as amino acid analogs having variant side chains with appropriate functional groups). For instance, the subject compound can include an amino acid analog such as, for example, cyanoalanine, canavanine, djenkolic acid, norleucine, 3-phosphoserine, homoserine, dihydroxy-phenylalanine, 5-hydroxytryptophan, 1-methylhistidine, 3-methylhistidine, diaminopimelic acid, ornithine, or
30 diaminobutyric acid. Other naturally occurring amino acid metabolites or precursors

having side chains which are suitable herein will be recognized by those skilled in the art and are included in the scope of the present invention.

Also included are the (D) and (L) stereoisomers of such amino acids when the structure of the amino acid admits of stereoisomeric forms. The configuration of the amino acids and amino acid residues herein are designated by the appropriate symbols (D), (L) or (DL), furthermore when the configuration is not designated the amino acid or residue can have the configuration (D), (L) or (DL). It will be noted that the structure of some of the compounds of this invention includes asymmetric carbon atoms. It is to be understood accordingly that the isomers arising from such asymmetry are included within the scope of this invention. Such isomers can be obtained in substantially pure form by classical separation techniques and by sterically controlled synthesis. For the purposes of this application, unless expressly noted to the contrary, a named amino acid shall be construed to include both the (D) or (L) stereoisomers.

The phrase "protecting group" as used herein means substituents which protect the reactive functional group from undesirable chemical reactions. Examples of such protecting groups include esters of carboxylic acids and boronic acids, ethers of alcohols and acetals and ketals of aldehydes and ketones. For instance, the phrase "N-terminal protecting group" or "amino-protecting group" as used herein refers to various amino-protecting groups which can be employed to protect the N-terminus of an amino acid or peptide against undesirable reactions during synthetic procedures. Examples of suitable groups include acyl protecting groups such as, to illustrate, formyl, dansyl, acetyl, benzoyl, trifluoroacetyl, succinyl and methoxysuccinyl; aromatic urethane protecting groups as, for example, benzyloxycarbonyl (Cbz); and aliphatic urethane protecting groups such as t-butoxycarbonyl (Boc) or 9-Fluorenylmethoxycarbonyl (Fmoc).

As noted above, certain compounds of the present invention may exist in particular geometric or stereoisomeric forms. The present invention contemplates all such compounds, including cis- and trans-isomers, R- and S-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric

carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

If, for instance, a particular enantiomer of a compound of the present invention is desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 67th Ed., 1986-87, inside cover. Also for purposes of this invention, the term "hydrocarbon" is contemplated to include all permissible compounds having at least one hydrogen and one carbon atom. In a broad aspect, the permissible hydrocarbons include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic organic compounds which can be substituted or unsubstituted.

A compound is said to have an "insulinotropic activity" if it is able to stimulate, or cause the stimulation of, the synthesis or expression of the hormone insulin.

The phrase "targeted by" refers to cells which are attacked and/or gradually destroyed by an autoimmune disease. For example, in Type I diabetes white blood cells called T lymphocytes produce immune factors called cytokines that attack and gradually destroy the beta cells of the pancreas.

iii. Exemplary Formulations

The inhibitors useful in the subject methods possess, in certain embodiments, the ability to lower blood glucose levels, to relieve obesity, to alleviate impaired

glucose tolerance, to inhibit hepatic glucose neogenesis, to inhibit diabetic ketoacidosis, and to lower blood lipid levels and to inhibit aldose reductase. They are thus useful for the prevention and/or therapy of hyperglycemia, obesity, hyperlipidemia, diabetic complications (including retinopathy, nephropathy, neuropathy, cataracts, coronary artery disease and arteriosclerosis) and furthermore for obesity-related hypertension and osteoporosis.

Diabetes mellitus is a disease characterized by hyperglycemia occurring from a relative or absolute decrease in insulin secretion, decreased insulin sensitivity or insulin resistance. The morbidity and mortality of this disease result from vascular, renal, and neurological complications. An oral glucose tolerance test is a clinical test used to diagnose diabetes. In an oral glucose tolerance test, a patient's physiological response to a glucose load or challenge is evaluated. After ingesting the glucose, the patient's physiological response to the glucose challenge is evaluated. Generally, this is accomplished by determining the patient's blood glucose levels (the concentration of glucose in the patient's plasma, serum or whole blood) for several predetermined points in time.

As described in the appended examples, we demonstrate that, *in vivo*, high affinity inhibitors of DPIV are biologically active with respect to regulation of glucose metabolism. For example, a single injection of the inhibitor Pro-boro-Pro (see examples for structure) was alone sufficient to improve glucose control. A single injection of Pro-boro-Pro was also observed to potentiate the response to a subtherapeutic dose of GLP-1. We have also observed that chronic (>5 days) treatment with Pro-boro-Pro alone lowers both fasting blood sugars, and the glycemic excursion to oral glucose challenge.


As indicated above, the inhibitors useful in the subject method can be peptide- or peptidomimetic-derived inhibitors of the target proteolytic activity, or can be a non-peptide compound identified, e.g., by drug screening assays described herein.

As discussed further below, a variety of assays are available in the art for identifying potential inhibitors of DPIV and the like, as well as assessing the various biological activities (including side-effects and toxicity) of such an inhibitor.

A. Examples of peptidyl DPIV inhibitors

In the case of DPIV inhibitors, a preferred class of inhibitors are peptidyl compounds based on the dipeptides Pro-Pro or Ala-Pro. Another preferred class of peptidyl inhibitors are compounds based on the dipeptide (D)-Ala-(L)-Ala. In many embodiments, it will be desirable to provide the peptidyl moiety as a peptidomimetic, e.g., to increase bioavailability and/or increase the serum half-life relative to the equivalent peptide. For instance, a variety of peptide backbone analogs are available in the art and be readily adapted for use in the subject methods.


In an exemplary embodiment, the peptidomimetic can be derived as a retro-inverso analog of the peptide. To illustrate, certain of the subject peptides can be generated as the retro-inverso analog (shown in its unprotected state):



Such retro-inverso analogs can be made according to the methods known in the art, such as that described by the Sisto et al. U.S. Patent 4,522,752. For example, the illustrated retro-inverso analog can be generated as follows. The geminal diamine corresponding to the N-terminal amino acid analogs is synthesized by treating an N-Boc-protected amino acid (having the sidechain R) with ammonia under HOBT-DCC coupling conditions to yield amide, and then effecting a Hofmann-type rearrangement with 1,1-bis-(trifluoroacetoxy)iodobenzene (TIB), as described in Radhakrishna et al. (1979) *J. Org. Chem.* 44:1746. The product amine salt is then coupled to a side-chain protected (e.g., as the benzyl ester) N-Fmoc D-enantiomer of the second amino acid residue (e.g., having a sidechain R') under standard conditions to yield the pseudodipeptide. The Fmoc (fluorenylmethoxycarbonyl) group is removed with piperidine in dimethylformamide, and the resulting amine is trimethylsilylated with bistrimethylsilylacetamide (BSA) before condensation with suitably alkylated, side-chain protected derivative of Meldrum's acid, as described in U.S. Patent 5,061,811 to


Pinori et al., to yield the retro-inverso tripeptide analog. The pseudotripeptide is then coupled with (protected) boro-proline under standard conditions to give the protected tetrapeptide analog. The protecting groups are removed to release the final product, which is purified by HPLC.

- 5 In another illustrative embodiment, the peptidomimetic can be derived as a retro-enantio analog of the peptide:




- 10 Retro-enantio analogs such as this can be synthesized using D-enantiomers of commercially available D-amino acids or other amino acid analogs and standard solid- or solution-phase peptide-synthesis techniques.

In still another illustrative embodiment, trans-olefin derivatives can be made with the subject boronophenylalanine analogs. For example, an exemplary olefin analog is:



- 15 The trans olefin analog can be synthesized according to the method of Y.K. Shue et al. (1987) *Tetrahedron Letters* 28:3225.

Still another class of peptidomimetic boronophenylalanine derivatives include the phosphonate derivatives, such as:



The synthesis of such phosphonate derivatives can be adapted from known synthesis schemes. See, for example, Loots et al. in *Peptides: Chemistry and Biology*, (Escom Science Publishers, Leiden, 1988, p. 118); Petrillo et al. in *Peptides: Structure and Function* (Proceedings of the 9th American Peptide Symposium, Pierce Chemical Co. Rockland, IL, 1985).

B. Non-peptidyl DPIP inhibitors

The pharmaceutical industry has developed a variety of different strategies for assessing millions of compounds a year as potential lead compounds based on inhibitory activity against an enzyme. DPIP and other proteolytic enzymes targeted by the subject method are amenable to the types of high throughput screening required to sample large arrays of compounds and natural extracts for suitable inhibitors.

As an illustrative embodiment, the ability of a test agent to inhibit DPIP can be assessed using a colorimetric or fluorometric substrate, such as Ala-Pro-paranitroanilide. See US Patent 5,462,928. Moreover, DPIP can be purified, and is accordingly readily amenable for use in such high throughput formats as multi-well plates.

Briefly, DPIP is purified from pig kidney cortex (Barth et al. (1974) Acta Biol Med Germ 32:157; Wolf et al. (1972) Acta Bio Mes Germ 37:409) or human placenta (Puschel et al. (1982) Eur J Biochem 126:359). An illustrative reaction mixture includes 50µM sodium Hepes (pH7.8), 10µM Ala-Pro-paranitroanilide, 6 milliunits of DPIP, and 2% (v/v) dimethylformamide in a total volume of 1.0 mL. The reaction is initiated by addition of enzyme, and formation of reaction product (paranitroanilide) in the presence and absence of a test compound can be detected photometrically, e.g., at 410 nm.

Exemplary compounds which can be screened for activity against DPIP (or other relevant enzymes) include peptides, nucleic acids, carbohydrates, small organic

molecules, and natural product extract libraries, such as isolated from animals, plants, fungus and/or microbes.

C. Assays of Insulinotropic Activity

5 In selecting a compound suitable for use in the subject method, it is noted that the insulinotropic property of a compound may be determined by providing that compound to animal cells, or injecting that compound into animals and monitoring the release of immunoreactive insulin (IRI) into the media or circulatory system of the animal, respectively. The presence of IRI can be detected through the use of a
10 radioimmunoassay which can specifically detect insulin.

Nonobese diabetic (NOD) mice are a well established model of type I diabetes (IDDM). In most litters, a prediabetic (>20 weeks) phase is observed which is characterized by pancreatic insulinitis without hyperglycemia. The NOD mice can purchased from, for example, The Jackson Laboratories (Bar Harbor, Me.). In an
15 exemplary embodiment, for treatment of the mice with a regimen including a DPIV inhibitor or control, sub-orbital sinus blood samples are taken before and at some time (e.g., 60 minutes) after dosing of each animal. Blood glucose measurements can be made by any of several conventional techniques, such as using a glucose meter. The blood glucose levels of the control and DPIV inhibitor dosed animals are compared

20 The metabolic fate of exogenous GLP-1 can also be followed in either nondiabetic and type I diabetic subjects, and the effect of a candidate DPIV inhibitor determined. For instance, a combination of high-pressure liquid chromatography (HPLC), specific radioimmunoassays (RIAs), and a enzyme-linked immunosorbent assay (ELISA), can be used, whereby intact biologically active GLP-1 and its
25 metabolites can be detected. See, for example, Deacon et al. (1995) Diabetes 44:1126-1131. To illustrate, after GLP-1 administration, the intact peptide can be measured using an NH₂-terminally directed RIA or ELISA, while the difference in concentration between these assays and a COOH-terminal-specific RIA allowed determination of NH₂-terminally truncated metabolites. Without inhibitor,
30 subcutaneous GLP-1 is rapidly degraded in a time-dependent manner, forming a

metabolite which co-elutes on HPLC with GLP-I(9-36) amide and has the same immunoreactive profile. For instance, thirty minutes after subcutaneous GLP-1 administration to diabetic patients ($n = 8$), the metabolite accounted for $88.5 \pm 1.9\%$ of the increase in plasma immunoreactivity determined by the COOH-terminal RIA, which was higher than the levels measured in healthy subjects ($78.4 \pm 3.2\%$; $n = 8$; $P < 0.05$). See Deacon et al., *supra*. Intravenously infused GLP-I was also extensively degraded.

D. Pharmaceutical Formulations

The inhibitors can be administered in various forms, depending on the disorder to be treated and the age, condition and body weight of the patient, as is well known in the art. For example, where the compounds are to be administered orally, they may be formulated as tablets, capsules, granules, powders or syrups; or for parenteral administration, they may be formulated as injections (intravenous, intramuscular or subcutaneous), drop infusion preparations or suppositories. For application by the ophthalmic mucous membrane route, they may be formulated as eyedrops or eye ointments. These formulations can be prepared by conventional means, and, if desired, the active ingredient may be mixed with any conventional additive, such as an excipient, a binder, a disintegrating agent, a lubricant, a corrigent, a solubilizing agent, a suspension aid, an emulsifying agent or a coating agent. Although the dosage will vary depending on the symptoms, age and body weight of the patient, the nature and severity of the disorder to be treated or prevented, the route of administration and the form of the drug, in general, a daily dosage of from 0.01 to 2000 mg of the compound is recommended for an adult human patient, and this may be administered in a single dose or in divided doses.

Glucose metabolism can be altered, and symptoms associated with type I diabetes can be decreased or eliminated, in accordance with a "timed" administration of DPPIV inhibitors wherein one or more appropriate indices for glucose metabolism and/or type I diabetes can be used to assess effectiveness of the treatment (dosage

and/or timing): e.g. glucose tolerance, glucose level, insulin level, insulin sensitivity, glycosylated hemoglobin.

An effective time for administering DPIV inhibitors needs to be identified. This can be accomplished by routine experiment as described below, using one or
5 more groups of animals (preferably at least 5 animals per group).

In animals, insulintropic activity by DPIV inhibitor treatment can be assessed by administering the inhibitor at a particular time of day and measuring the effect of the administration (if any) by measuring one or more indices associated with glucose metabolism, and comparing the post-treatment values of these indices to the
10 values of the same indices prior to treatment.

The precise time of administration and/or amount of DPIV inhibitor that will yield the most effective results in terms of efficacy of treatment in a given patient will depend upon the activity, pharmacokinetics, and bioavailability of a particular compound, physiological condition of the patient (including age, sex, disease type and
15 stage, general physical condition, responsiveness to a given dosage and type of medication), route of administration, etc. However, the above guidelines can be used as the basis for fine-tuning the treatment, e.g., determining the optimum time and/or amount of administration, which will require no more than routine experimentation consisting of monitoring the subject and adjusting the dosage and/or timing.

20 While the subject is being treated, glucose metabolism is monitored by measuring one or more of the relevant indices at predetermined times during a 24-hour period. Treatment (amounts, times of administration and type of medication) may be adjusted (optimized) according to the results of such monitoring. The patient is periodically reevaluated to determine extent of improvement by measuring the
25 same parameters, the first such reevaluation typically occurring at the end of four weeks from the onset of therapy, and subsequent reevaluations occurring every 4 to 8 weeks during therapy and then every 3 months thereafter. Therapy may continue for several months or even years with six months being a typical length of therapy for humans.

Adjustments to the amount(s) of drug(s) administered and possibly to the time of administration may be made based on these reevaluations. For example, if after 4 weeks of treatment one of the metabolic indices has not improved but at least one other one has, the dose could be increased by 1/3 without changing the time of administration.

Treatment can be initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage should be increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

The phrase "therapeutically effective amount" as used herein means that amount of, e.g., a DPIV inhibitor(s), which is effective for producing some desired therapeutic effect by inhibiting, for example, the proteolysis of a peptide hormone at a reasonable benefit/risk ratio applicable to any medical treatment.

The phrase "pharmaceutically acceptable" is employed herein to refer to those DPIV inhibitors, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "pharmaceutically acceptable carrier" as used herein means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject chemical from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and

suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

The term "pharmaceutically acceptable salts" refers to the relatively non-toxic, inorganic and organic acid addition salts of DPIV inhibitors. These salts can be prepared in situ during the final isolation and purification of the DPIV Inhibitors, or by separately reacting a purified DPIV inhibitor in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and the like. (See, for example, Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19)

In other cases, the DPIV inhibitor useful in the methods of the present invention may contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable bases. The term "pharmaceutically-acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of a DPIV inhibitor(s). These salts can likewise be prepared in situ during the final isolation and purification of the DPIV inhibitor(s), or by separately reacting the purified DPIV inhibitor(s) in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation, with ammonia, or with a pharmaceutically-acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine,

ethanolamine, diethanolamine, piperazine and the like (see, for example, Berge et al., *supra*).

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, 5 sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as 10 ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Formulations useful in the methods of the present invention include those 15 suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal, aerosol and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the host 20 being treated, the particular mode of administration. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, out of one hundred per cent, this amount will range from about 1 per cent to about ninety-nine percent of active ingredient, preferably from about 5 per cent to 25 about 70 per cent, most preferably from about 10 per cent to about 30 per cent.

Methods of preparing these formulations or compositions include the step of bringing into association a DPIV inhibitor(s) with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a DPIV inhibitor with liquid carriers, or 30 finely divided solid carriers, or both, and then, if necessary, shaping the product.

Formulations suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or
5 syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a DPIV inhibitor(s) as an active ingredient. A compound may also be administered as a bolus, electuary or paste.

In solid dosage forms for oral administration (capsules, tablets, pills, dragees,
10 powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or
15 acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, acetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite
20 clay; (9) lubricants, such a talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as
25 lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent,
30 preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets

may be made by molding in a suitable machine a mixture of the powdered peptide or peptidomimetic moistened with an inert liquid diluent.

Tablets, and other solid dosage forms, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active DPIV inhibitor(s) may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

- 5 Formulations for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more DPIV inhibitor(s) with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in
10 the rectum or vaginal cavity and release the active agent.

Formulations which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

- 15 Dosage forms for the topical or transdermal administration of a DPIV inhibitor(s) include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active component may be mixed under sterile conditions with a pharmaceutically-acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

- 20 The ointments, pastes, creams and gels may contain, in addition to DPIV inhibitor(s), excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

- 25 Powders and sprays can contain, in addition to a DPIV inhibitor(s), excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

The DPIV inhibitor(s) can be alternatively administered by aerosol. This is accomplished by preparing an aqueous aerosol, liposomal preparation or solid

particles containing the compound. A nonaqueous (e.g., fluorocarbon propellant) suspension could be used. Sonic nebulizers are preferred because they minimize exposing the agent to shear, which can result in degradation of the compound.

Ordinarily, an aqueous aerosol is made by formulating an aqueous solution or
5 suspension of the agent together with conventional pharmaceutically acceptable carriers and stabilizers. The carriers and stabilizers vary with the requirements of the particular compound, but typically include nonionic surfactants (Tweens, Pluronic, or polyethylene glycol), innocuous proteins like serum albumin, sorbitan esters, oleic acid, lecithin, amino acids such as glycine, buffers, salts, sugars or sugar alcohols.
10 Aerosols generally are prepared from isotonic solutions.

Transdermal patches have the added advantage of providing controlled delivery of a DPIV inhibitor(s) to the body. Such dosage forms can be made by dissolving or dispersing the agent in the proper medium. Absorption enhancers can also be used to increase the flux of the peptidomimetic across the skin. The rate of
15 such flux can be controlled by either providing a rate controlling membrane or dispersing the peptidomimetic in a polymer matrix or gel.

Ophthalmic formulations, eye ointments, powders, solutions and the like, are also contemplated as being within the scope of this invention.

Pharmaceutical compositions of this invention suitable for parenteral
20 administration comprise one or more DPIV inhibitor(s) in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with
25 the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic

esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

5 These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the
10 injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material
15 having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of DPIV
20 inhibitor(s) in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are
25 compatible with body tissue.

When the DPIV inhibitor(s) of the present invention are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99.5% (more preferably, 0.5 to 90%) of active ingredient in combination with a pharmaceutically acceptable
30 carrier.

The preparations of agents may be given orally, parenterally, topically, or rectally. They are of course given by forms suitable for each administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc. administration by injection, infusion or inhalation; 5 topical by lotion or ointment; and rectal by suppositories. Oral administration is preferred.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, 10 intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the 15 administration of a DPIV inhibitor, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

These DPIV inhibitor(s) may be administered to humans and other animals for therapy by any suitable route of administration, including orally, nasally, as by, for 20 example, a spray, rectally, intravaginally, parenterally, intracisternally and topically, as by powders, ointments or drops, including buccally and sublingually.

Regardless of the route of administration selected, the DPIV inhibitor(s), which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically- 25 acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a

particular patient, composition, and mode of administration, without being toxic to the patient.

E. Conjoint administration

- 5 Another aspect of the invention provides a conjoint therapy wherein one or more other therapeutic agents are administered with the protease inhibitor. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of the treatment.

- 10 In one embodiment, a DPIV inhibitor is conjointly administered with immunosuppressive agents, such as, for example, cyclosporine; cyclosporine in conjunction with either azathioprine, steroids, or both; FK506 tacrolimus (Prograf); or mycophenolate mofetil (Cellcept).

iv. Business Methods

- 15 One aspect of the present invention relates to a kit comprising compounds as described herein, such as DPIV inhibitors, for treatment or prevention of autoimmune disorders, such as Type 1 diabetes, septic shock, multiple sclerosis, IBD or Crohn's disease in a patient, preferably a human, and in association with instructions (written and/or pictorial) describing the use of the formulation for treatment or prevention of
20 autoimmune disorders, such as Type 1 diabetes, septic shock, multiple sclerosis, IBD or Crohn's disease, and optionally, warnings of possible side effects and drug-drug or drug-food interactions.

- The invention further contemplates a method for conducting a pharmaceutical business, comprising: (a) manufacturing a pharmaceutical preparation comprising a
25 sterile pharmaceutical excipient and compounds as described herein, such as DPIV inhibitors; and (b) marketing (e.g., providing promotional and/or informative presentations (such as displays, telemarketing, and lectures), products (such as trial samples of the preparation), and/or documentation (including leaflets, pamphlets, websites, posters, etc.)) to healthcare providers, such as doctors, hospitals, clinics,

etc., a benefit of using the pharmaceutical preparation for treatment or prevention of autoimmune disorders, such as Type 1 diabetes, septic shock, multiple sclerosis, IBD or Crohn's disease.

5 Another aspect of the invention provides for a method for conducting a pharmaceutical business, comprising: (a) providing a distribution network for selling the pharmaceutical composition comprising a sterile pharmaceutical excipient and compounds as described herein, such as DPIV Inhibitors; and (b) providing instruction material to patients or physicians for using the pharmaceutical composition for treatment or prevention of autoimmune disorders, such as Type 1 diabetes, septic
10 shock, multiple sclerosis, IBD or Crohn's disease.

Yet another aspect of the invention provides for a method for conducting a pharmaceutical business, comprising: (a) determining an appropriate pharmaceutical preparation and dosage of a compounds as described herein, such as DPIV inhibitors for treatment or prevention of autoimmune disorders, such as Type 1 diabetes, septic
15 shock, multiple sclerosis, IBD or Crohn's disease; (b) conducting therapeutic profiling of the pharmaceutical preparation for efficacy and toxicity in animals; (c) providing a distribution network for selling a pharmaceutical composition having an acceptable therapeutic profile; and, optionally, (d) providing a sales group for marketing the preparation to healthcare providers.

20

Exemplification

The invention now being generally described, it will be more readily understood by reference to the following examples which are included merely for purposes of illustration of certain aspects and embodiments of the present invention,
25 and are not intended to limit the invention.

Example 1: Synthesis of BoroProline

Referring to Figure 1, the starting compound I is prepared essentially by the procedure of Matteson et al. (Organometallics 3:1284, 1984), except that a pinacol ester is substituted for the pinanediol ester. Similar compounds such as boropipecolic

acid and 2-azetidine boronic acid can be prepared by making the appropriate selection of starting material to yield the pentyl and propyl analogs of compound I. Further, Cl can be substituted for Br in the formula, and other diol protecting groups can be substituted for pinacol in the formula, e.g., 2, 3-butanediol and
5 alphapinanediol.

Compound II is prepared by reacting compound I with $[(CH_3)_3O_3Si]_2N-Li^+$. In this reaction hexamethyldisilazane is dissolved in tetrahydrofuran and an equivalent of n-butyllithium added at $-78^\circ C$. After warming to room temperature ($20^\circ C$) and cooling to $-78^\circ C$, an equivalent of compound I is added in tetrahydrofuran. The
10 mixture is allowed to slowly come to room temperature and to stir overnight. The alpha-bis[trimethylsilane]-protected amine is isolated by evaporating solvent and adding hexane under anhydrous conditions. Insoluble residue is removed by filtration under a nitrogen blanket, yielding a hexane solution of compound II.

Compound III, the N-trimethylsilyl protected form of boroProline is obtained
15 by the thermal cyclization of compound II during the distillation process in which compound II is heated to $100-150^\circ C$ and distillate is collected which boils $66-62^\circ C$ at $0.06-0.10$ mm pressure.

Compound IV, boroProline-pinacol hydrogen chloride, is obtained by treatment of compound III with HCl:dioxane. Excess HCl and by-products are
20 removed by trituration with ether. The final product is obtained in a high degree of purity by recrystallization from ethyl acetate.

The boroProline esters can also be obtained by treatment of the reaction mixture obtained in the preparation of compound II with anhydrous acid to yield 1-amino-4-bromobutyl boronate pinacol as a salt. Cyclization occurs after neutralizing
25 the salt with base and heating the reaction.

Example 2: Preparation of boroProline-pinacol

The intermediate, 4-Bromo-1-chlorobutyl boronate pinacol, was prepared by the method in Matteson et al. (Organometallics 3:1284, 1984) except that conditions were modified for large scale preparations and pinacol was substituted for the pinanediol protecting group.

3-bromopropyl boronate pinacol was prepared by hydrogenboronation of allyl bromide (173 ml, 2.00 moles) with catechol borane (240 ml, 2.00 moles). Catechol borane was added to allyl bromide and the reaction heated for 4 hours at 100°C under a nitrogen atmosphere. The product, 3-bromopropyl boronate catechol (bp 95-102°C, 0.25 mm), was isolated in a yield of 49% by distillation. The catechol ester (124 g, 0.52 moles) was transesterified with pinacol (61.5 g, 0.52 moles) by mixing the component in 50 ml of THF and allowing them to stir for 0.5 hours at 0°C and 0.5 hours at room temperature. Solvent was removed by evaporation and 250 ml of hexane added. Catechol was removed as a crystalline solid. Quantitative removal was achieved by successive dilution to 500 ml and to 1000 ml with hexane and removing crystals at each dilution. Hexane was evaporated and the product distilled to yield 177g (bp 60-64°C, 0.35 mm).

4-Bromo-1-chlorobutyl boronate pinacol was prepared by homologation of the corresponding propyl boronate. Methylene chloride (50.54 ml, 0.713 moles) was dissolved in 500 ml of THF, 1.54N n-butyllithium in hexane (480 ml, 0.780 moles) was slowly added at -100°C. 3-Bromopropyl boronate pinacol (178 g, 0.713 moles) was dissolved in 500 ml of THG, cooled to the freezing point of the solution, and added to the reaction mixture. Zinc chloride (54.4 g, 0.392 moles) was dissolved in 250 ml of THG, cooled to 0°C, and added to the reaction mixture in several portions. The reaction was allowed to slowly warm to room temperature and to stir overnight. Solvent was evaporated and the residue dissolved in hexane (1 liter) and washed with water (1 liter). Insoluble material was discarded. After drying over anhydrous magnesium sulfate and filtering, solvent was evaporated. The product was distilled to yield 147 g (bp 110-112°C, 0.200 mm).

N-Trimethylsilyl-boroProline pinacol was prepared first by dissolving hexamethyldisilazane (20.0 g, 80.0 mmoles) in 30 ml of THF, cooling the solution to -78°C, and adding 1.62N n-butyllithium in hexane (49.4 ml, 80.0 mmoles). The solution was allowed to slowly warm to room temperature. It was recooled to -78°C.
5 and 4-bromo-1-chlorobutyl boronate pinacol (23.9 g, 80.0 mmoles) added in 20 ml of THF. The mixture was allowed to slowly warm to room temperature and to stir overnight. Solvent was removed by evaporation and dry hexane (400 ml) added to yield a precipitant which was removed by filtration under a nitrogen atmosphere. The filtrate was evaporated and the residue distilled, yielding 19.4 g of the desired product
10 (bp 60-62°C, 0.1-0.06 mm).

H-boroProline-pinacol.HCl (boroProline-pinacol.HCl) was prepared by cooling N-trimethylsilyl-boroProline pinacol (16.0 g, 61.7 mmoles) to -78°C and adding 4N HCL:dioxane 46 ml, 185 mmoles). The mixture was stirred 30 minutes at -78°C and 1 hour at room temperature. Solvent was evaporated and the residue
15 triturated with ether to yield a solid. The crude product was dissolved in chloroform and insoluble material removed by filtration. The solution was evaporated and the product crystallized from ethyl acetate to yield 11.1 g of the desired product (mp 156.5-157°C).

20 *Example 3: Synthesis of boroProline Peptides*

General methods of coupling of N-protected peptides and amino acids with suitable side-chain protecting groups to H-boroProline-pinacol are applicable. When needed, side-chain protecting and N-terminal protecting groups can be removed by treatment with anhydrous HCl, HBr, trifluoroacetic acid, or by catalytic
25 hydrogenation. These procedures are known to those skilled in the art of peptide synthesis.

The mixed anhydride procedure of Anderson et al. (J. Am. Chem. Soc. 89:5012, 1984) is preferred for peptide coupling. Referring again to Figure 1, the mixed anhydride of an N-protected amino acid or a peptide is prepared by dissolving
30 the peptide in tetrahydrofuran and adding one equivalent of N-methylmorpholine. The

solution is cooled to -20°C and an equivalent of isobutyl chloroformate is added. After 5 minutes, this mixture and one equivalent of triethylamine (or other sterically hindered base) are added to a solution of H-boroPro-pinacol dissolved in either cold chloroform or tetrahydrofuran.

- 5 The reaction mixture is routinely stirred for one hour at -20°C and 1 to 2 hours at room temperature (20°C). Solvent is removed by evaporation, and the residue is dissolved in ethyl acetate. The organic solution is washed with 0.20N hydrochloric acid, 5% aqueous sodium bicarbonate, and saturated aqueous sodium chloride. The organic phase is dried over anhydrous sodium sulfate, filtered, and evaporated.
- 10 Products are purified by either silica gel chromatography or gel permeation chromatography using Sephadex TM LH-20 and methanol as a solvent.

- Previous studies have shown that the pinacol protecting group can be removed in situ by preincubation in phosphate buffer prior to running biological experiments (Kettner et al., J. Biol. Chem. 259:15106, 1984). Several other methods are also
- 15 applicable for removing pinacol groups from peptides, including boroProline, and characterizing the final product. First, the peptide can be treated with diethanolamine to yield the corresponding diethanolamine boronic acid ester, which can be readily hydrolyzed by treatment with aqueous acid or a sulfonic acid substituted polystyrene resin as described in Kettner et al. (supra). Both pinacol and pinanediol protecting
- 20 groups can be removed by treating with BC13 in methylene chloride as described by Kinder et al. (J. Med. Chem. 28:1917). Finally, the free boronic acid can be converted to the difluoroboron derivative (-BF₂) by treatment with aqueous HF as described by Kinder et al. (supra).

- Similarly, different ester groups can be introduced by reacting the free boronic
- 25 acid with various di-hydroxy compounds (for example, those containing heteroatoms such as S or N) in an inert solvent.

Example 4: Preparation of H-Ala-boroPro

Boc-Ala-boroPro was prepared by mixed anhydride coupling of the N-Boc-protected alanine and H-boroPro prepared as described above. H-Ala-boroPro (Ala-boroPro) was prepared by removal of the Boc protecting group at 0°C in 3.5 molar excess of 4N HCl-dioxane. The coupling and deblocking reactions were performed by standard chemical reaction. Ala-boroPro has a K_i for DP-IV of in the nanomolar range. Boc-blocked Ala-boroPro has no affinity for DP-IV.

The two diastereomers of Ala-boroPro-pinacol, L-Ala-D-boroPro-pinacol and L-Ala-L-boroPro-pinacol, can be partially separated by silica gel chromatography with 20% methanol in ethyl acetate as eluant. The early fraction appears by NMR analysis to be 95% enriched in one isomer. Because this fraction has more inhibits DP-IV to a greater extent than later fractions (at equal concentrations) it is probably enriched in the L-boroPro (L-Ala-L-boroPro-pinacol) isomer.

Example 5: Synthesis of Cyclohexylglycine boroAla

Referring to Figure 1, a solution of 515 mg (2.00 mmol) of Boc-L-2-(cyclohexyl)glycine 1 (Chem-Impex International), 587 mg (2.26 mmol) of HCl.boroAla pinane 2, 332 mg (2.46 mmol) of HOBT, and 671 μ L (4.84 mmol) of triethylamine in 6 mL of anhydrous DMF was treated with 498 mg (2.60 mmol) of EDC, and the resulting solution stirred at room temperature under argon for 18 h. The reaction mixture was diluted with a 200 mL of 10% aqueous citric acid and the resulting mixture extracted with 2 x 100 mL of ethyl acetate. The combined extracts were washed with brine, dried (MgSO_4), filtered, and concentrated to give a clear oil. The crude oil was chromatographed over silica gel with ethyl acetate/hexane to give the product ester as a clear oil. The oil was then dissolved in hydrogen chloride in diethyl ether (1.0 M solution, 25 mL) and stirred for 48 hours at room temperature. The mixture was evaporated to dryness *in vacuo* and redissolved in 25 mL phenylboronic acid solution (244 mg, 2 mmol) at pH 2 (0.01 N HCl) and ether (25 mL). After stirring for 30 min, the ether layer was removed and replaced with fresh ether (25 mL). This step was repeated for four times. The aqueous phase was then lyophilized and purified by HPLC to afford 170 mg (37%) of the target compound 3.

Example 6: DPIV Assays on Serum Samples from Rats

Experiments show that DPIV enzyme activity was significantly decreased in rats treated with Cyclohexylglycine-boroAla. See Figure 7. Four rats were used in this experiment: two females (#3 and #9) and two males (#10 and #11). Blood and plasma samples were collected from rats 1 hour after being treated with Cyclohexylglycine-boroAla. The collected serum samples were evaluated for DPPIV activity of Cyclohexylglycine-boroAla as follows:

10

1. 2 mg of Ala-Pro-paranitroanalide (substrate) was dissolved in 20 ml 0.1 M HEPES pH 8, 0.14 M NaCl (buffer).
2. Serum samples were diluted into substrate solution in the wells of a microtiter plate. For each sample, 10uL of serum was diluted into 150 μ L of substrate.
- 15 3. A reading of the A410 in each well was recorded immediately after the dilution of serum into substrate, and again after approximately 1 hour. The time of data acquisition for each reading is recorded in the data file by the microplate reader software.

The rate of absorbance change was obtained by subtracting the first reading from the second and dividing by the reaction time to give DeltaA410/hr. The DPIV activity was plotted in units of DeltaA410 $\text{hr}^{-1} \mu\text{L}^{-1}$.

20

Example 7: Incidence of Diabetes in NOD mice upon treatment with DPIV Inhibitors

The experiment was started on 8-10 week old NOD mice. The mice were kept under VAF/SPF conditions and fed everyday with either Val-boro-Pro (0.034 mg/kg) or Cyclohexyl-boro-Ala (0.34 mg/kg) for 60 days and then observed for the development of spontaneous diabetes. The mice were tested for the excretion of sugar in urine and considered positive for diabetes when sugar was detected in the urine.

25

The graph shows cumulative incidence of Diabetes over time (Figure 5). One of the interesting features of the NOD mice treated with Val-boro-Pro was that some of the mice that showed signs of diabetes got better and recovered. At the end of the experiment, the mice in Val-boro-Pro group not only had low incidence of Diabetes
5 (Figure 6) but also showed generally good health compared to the other groups.

All of the above-cited references and publications are hereby incorporated by reference.


Equivalents

10 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

Claims:

1. A method for treating an autoimmune disease in an animal involving tissue which is responsive to a peptide hormone processed by dipeptidylpeptidase IV (DPIV), comprising administering to said animal a composition comprising one or more inhibitors of DPIV in an amount sufficient to promote tissue regeneration of tissue targeted by the autoimmune disease.
2. The method of claim 1, wherein the autoimmune disease is selected from: Type I diabetes, septic shock, multiple sclerosis, inflammatory bowel disease (IBD) and Crohn's disease.
3. The method of claim 1, wherein the peptide hormone is selected from: glucagon, glucagon-like peptide-1 (GLP-1), glucagon-like peptide-2 (GLP-2, PG 126-158), glicentin (corresponding to PG 1-69), oxyntomodulin (PG 33-69), glicentin-related pancreatic polypeptide (GRPP, PG 1-30), intervening peptide-2 (IP-2, PG 111-122amide), glucose-dependent insulintropic peptides, vasoactive intestinal polypeptide (VIP), vasostatin I or II, peptide histidine methionine (PHM), peptide histidine isoleucine (PHI), secretin, gastric inhibitory peptide, gastrin-releasing peptide (GRP), growth hormone-releasing hormone (GHRH), helospectin, helodermin, pituitary adenylate cyclase-activating peptide (PACAP, PACAP 27, and PACAP 38), and PACAP-related peptide (PRP), peptide YY (PYY), neuropeptide Y (NPY), gastric inhibitory peptide (GIP), helodermin, peptide histidine isoleucine (PHI), as well as calcitonin and secretin.
4. The method of claim 1, wherein the inhibitor is peptidomimetic of a peptide selected from the group consisting of Pro-Pro, Ala-Pro, and (D)-Ala-(L)-Ala.
5. The method of claim 1, wherein the inhibitor has a molecular weight less than 7500 amu.

6. The method of claim 1, wherein the inhibitor is orally active.
7. The method of claim 1, wherein the inhibitor is represented by the general formula:



5

wherein

A represents a 4-8-membered heterocycle including the N and the C α carbon;


Z represents C or N;

10

W represents a functional group which reacts with an active site residue of the targeted protease,

R₁ represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or

15




20

R₂ is absent or represents one or more substitutions to the ring A, each of which can independently be a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl, a thiocarbonyl, an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -lower alkyl, $-(CH_2)_m-O$ -lower alkenyl, $-(CH_2)_n-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -lower alkyl, $-(CH_2)_m-S$ -lower alkenyl, $-(CH_2)_n-S-(CH_2)_m-R_7$;

if X is N, R₃ represents hydrogen, if X is C, R₃ represents hydrogen or a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl, a thiocarbonyl, an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, -(CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-lower alkyl, -(CH₂)_m-O-lower alkenyl, -(CH₂)_n-O-(CH₂)_m-R₇, -(CH₂)_m-SH, -(CH₂)_m-S-lower alkyl, -(CH₂)_m-S-lower alkenyl, -(CH₂)_n-S-(CH₂)_m-R₇;

R₆ represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, -(CH₂)_m-R₇, -(CH₂)_m-OH, -(CH₂)_m-O-alkyl, -(CH₂)_m-O-alkenyl, -(CH₂)_m-O-alkynyl, -(CH₂)_m-O-(CH₂)_m-R₇, -(CH₂)_m-SH, -(CH₂)_m-S-alkyl, -(CH₂)_m-S-alkenyl, -(CH₂)_m-S-alkynyl, or -(CH₂)_m-S-(CH₂)_m-R₇,



R₇ represents, independently for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

m is zero or an integer in the range of 1 to 8; and


n is an integer in the range of 1 to 8.

8. The method of claim 7,

wherein

20

W represents -CN, -CH=NR₅,



5 R_5 represents H, an alkyl, an alkenyl, an alkynyl, $-\text{C}(\text{X}_1)(\text{X}_2)\text{X}_3$, $-(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_n\text{-OH}$, $-(\text{CH}_2)_n\text{-O-alkyl}$, $-(\text{CH}_2)_n\text{-O-alkenyl}$, $-(\text{CH}_2)_n\text{-O-alkynyl}$, $-(\text{CH}_2)_n\text{-O}-(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_n\text{-SH}$, $-(\text{CH}_2)_n\text{-S-alkyl}$, $-(\text{CH}_2)_n\text{-S-alkenyl}$, $-(\text{CH}_2)_n\text{-S-alkynyl}$, $-(\text{CH}_2)_n\text{-S}-(\text{CH}_2)_m\text{-R}_7$, $-\text{C}(\text{O})\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{C}(\text{O})\text{OR}'_7$;

R_7 represents, independently for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

10 R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle; and

Y_1 and Y_2 can, independently, be OH, or a group capable of being hydrolyzed to a hydroxyl group, including cyclic derivatives where Y_1 and Y_2 taken together form a ring having from 5 to 8 atoms in the ring structure,

15 R_{50} represents O or S;


R_{51} represents N_3 , SH_2 , NH_2 , NO_2 or OR'_7 ;

20 R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure

X_1 represents a halogen;


X_2 and X_3 each represent, independently, a hydrogen or a halogen.

9. The method of claim 8, wherein the ring A is represented by the formula




wherein n is an integer of 1 or 2.

10. The method of claim 8, wherein W represents



11. The method of claim 8, wherein R₁ represents




wherein

R₃₆ is a small hydrophobic group and R₃₈ is hydrogen, or, R₃₆ and R₃₈ together form a 4-7-membered heterocycle including the N and the C α carbon, as defined for A above; and

R₄₀ represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group.


12. The method of claim 8, wherein R₂ is absent, or represents a small hydrophobic group.

13. The method of claim 8, wherein R_3 is a hydrogen, or a small hydrophobic group.
14. The method of claim 8, wherein R_5 is a hydrogen, or a halogenated lower alkyl.
15. The method of claim 8, wherein X_1 is a fluorine, and X_2 and X_3 , if halogens, are fluorine.
16. The method of claim 1, wherein the inhibitor is represented by the general formula:




wherein

R_1 represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or



R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-alkynyl$, $-(CH_2)_m-S-(CH_2)_m-R_7$,



R_7 represents an aryl, a cycloalkyl, a cycloalkenyl, or a heterocycle;


R_8 and R_9 each, independently, represent hydrogen, alkyl, alkenyl, -
 $(\text{CH}_2)_m\text{---R}_7$, -C(=O)-alkyl, -C(=O)-alkenyl, -C(=O)-alkynyl, -C(=O)- $(\text{CH}_2)_m\text{---}$
 5 R_7 , or R_8 and R_9 taken together with the N atom to which they are attached
 complete a heterocyclic ring having from 4 to 8 atoms in the ring structure;

R_{11} and R_{12} each, independently, represent hydrogen, a alkyl, or a
 pharmaceutically acceptable salt, or R_{11} and R_{12} taken together with the O-B-
 O atoms to which they are attached complete a heterocyclic ring having from
 10 5 to 8 atoms in the ring structure;

m is zero or an integer in the range of 1 to 8; and


n is an integer in the range of 1 to 8.

17. The method of claim 1, wherein the inhibitor is represented by the general
 15 formula




wherein,

R_1 represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or



5 R_6 represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, $-(\text{CH}_2)_m-\text{R}_7$, $-(\text{CH}_2)_m-\text{OH}$, $-(\text{CH}_2)_m-\text{O-alkyl}$, $-(\text{CH}_2)_m-\text{O-alkenyl}$, $-(\text{CH}_2)_m-\text{O-alkynyl}$, $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_m-\text{R}_7$, $-(\text{CH}_2)_m-\text{SH}$, $-(\text{CH}_2)_m-\text{S-alkyl}$, $-(\text{CH}_2)_m-\text{S-alkenyl}$, $-(\text{CH}_2)_m-\text{S-alkynyl}$, $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_m-\text{R}_7$,



10

R_7 represents an aryl, a cycloalkyl, a cycloalkenyl, or a heterocycle;


R_8 and R_9 each independently represent hydrogen, alkyl, alkenyl, $-(\text{CH}_2)_m-\text{R}_7$, $-\text{C}(=\text{O})-\text{alkyl}$, $-\text{C}(=\text{O})-\text{alkenyl}$, $-\text{C}(=\text{O})-\text{alkynyl}$, $-\text{C}(=\text{O})-(\text{CH}_2)_m-\text{R}_7$, or R_8 and R_9 taken together with the N atom to which they are attached complete a heterocyclic ring having from 4 to 8 atoms in the ring structure; and

15

m is zero or an integer in the range of 1 to 8; and


n is an integer in the range of 1 to 8.

18. The method of claim 1, wherein the inhibitor is represented by the general formula:






wherein

- 5 R_1 represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or



- 10 R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-alkynyl$, $-(CH_2)_m-S-(CH_2)_m-R_7$,



- 15 R_7 represents, independently for each occurrence, an aryl, a cycloalkyl, a cycloalkenyl, or a heterocycle;


R_8 and R_9 each independently represent hydrogen, alkyl, alkenyl, $-(CH_2)_m-R_7$, $-C(=O)-alkyl$, $-C(=O)-alkenyl$, $-C(=O)-alkynyl$, $-C(=O)-(CH_2)_m-R_7$, or R_8 and R_9 taken together with the N atom to which they are attached complete a heterocyclic ring having from 4 to 8 atoms in the ring structure;

5 X_1 , X_2 and X_3 each represent a hydrogen or a halogen; and

m is zero or an integer in the range of 1 to 8; and

n is an integer in the range of 1 to 8.

10 19. The method of claim 1, wherein the inhibitor is represented by the general formula:




wherein

R_{32} is a small hydrophobic group; and


15 R_{30} represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group.

20 20. The method of claim 1, wherein the inhibitor is represented by the general formula:




wherein

W represents a functional group which reacts with an active site residue of the targeted protease, as for example, $-\text{CN}$, $-\text{CH}=\text{NR}_5$,




R_1 represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or



R_3 represents hydrogen or a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl, a thiocarbonyl, an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, $-(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-OH}$, $-(\text{CH}_2)_m\text{-O-lower alkyl}$, $-(\text{CH}_2)_m\text{-O-lower alkenyl}$, $-(\text{CH}_2)_n\text{-O-}(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-SH}$, $-(\text{CH}_2)_m\text{-S-lower alkyl}$, $-(\text{CH}_2)_m\text{-S-lower alkenyl}$, $-(\text{CH}_2)_n\text{-S-}(\text{CH}_2)_m\text{-R}_7$;

R_5 represents H, an alkyl, an alkenyl, an alkynyl, $-\text{C}(\text{X}_1)(\text{X}_2)\text{X}_3$, $-(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_n\text{-OH}$, $-(\text{CH}_2)_n\text{-O-alkyl}$, $-(\text{CH}_2)_n\text{-O-alkenyl}$, $-(\text{CH}_2)_n\text{-O-alkynyl}$, $-(\text{CH}_2)_n\text{-O-}(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_n\text{-SH}$, $-(\text{CH}_2)_n\text{-S-alkyl}$, $-(\text{CH}_2)_n\text{-S-alkenyl}$, $-(\text{CH}_2)_n\text{-S-alkynyl}$, $-(\text{CH}_2)_n\text{-S-}(\text{CH}_2)_m\text{-R}_7$, $-\text{C}(\text{O})\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{C}(\text{O})\text{OR}'_7$;

R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, $-(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-OH}$, $-(\text{CH}_2)_m\text{-O-alkyl}$, $-(\text{CH}_2)_m\text{-O-alkenyl}$, $-(\text{CH}_2)_m\text{-O-alkynyl}$, $-(\text{CH}_2)_m\text{-O-}(\text{CH}_2)_m\text{-R}_7$, $-(\text{CH}_2)_m\text{-SH}$, $-(\text{CH}_2)_m\text{-S-alkyl}$, $-(\text{CH}_2)_m\text{-S-alkenyl}$, $-(\text{CH}_2)_m\text{-S-alkynyl}$, $-(\text{CH}_2)_m\text{-S-}(\text{CH}_2)_m\text{-R}_7$,



R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

5 R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R_{61} and R_{62} , independently, represent small hydrophobic groups;

10 Y_1 and Y_2 can, independently, be OH, or a group capable of being hydrolyzed to a hydroxyl group, including cyclic derivatives where Y_1 and Y_2 taken together form a ring having from 5 to 8 atoms in the ring structure,

R_{50} represents O or S;

R_{51} represents N_3 , SH_2 , NH_2 , NO_2 or OR'_7 ;

15 R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure

X_1 represents a halogen;


X_2 and X_3 each, independently, represent a hydrogen or a halogen;

m is zero or an integer in the range of 1 to 8; and

n is an integer in the range of 1 to 8.

21. The method of claim 4, wherein the boronyl peptidomimetic is represented in the general formula:

5



wherein

each A independently represents a 4-8 membered heterocycle including the N and the C α carbon;

10

R_2 is absent or represents one or more substitutions to the ring A, each of which can independently be a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl, a thiocarbonyl, an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -lower alkyl, $-(CH_2)_m-O$ -lower alkenyl, $-(CH_2)_n-O$ - $(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -lower alkyl, $-(CH_2)_m-S$ -lower alkenyl, $-(CH_2)_n-S-(CH_2)_m-R_7$;

15

R_3 represents hydrogen or a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl, a thiocarbonyl, an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -lower alkyl, $-(CH_2)_m-O$ -lower alkenyl, $-(CH_2)_n-O$ - $(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -lower alkyl, $-(CH_2)_m-S$ -lower alkenyl, $-(CH_2)_n-S-(CH_2)_m-R_7$;


20

5

R_5 represents H, an alkyl, an alkenyl, an alkynyl, $-C(X_1)(X_2)X_3$, $-(CH_2)_m-R_7$, $-(CH_2)_n-OH$, $-(CH_2)_n-O-alkyl$, $-(CH_2)_n-O-alkenyl$, $-(CH_2)_n-O-alkynyl$, $-(CH_2)_n-O-(CH_2)_m-R_7$, $-(CH_2)_n-SH$, $-(CH_2)_n-S-alkyl$, $-(CH_2)_n-S-alkenyl$, $-(CH_2)_n-S-alkynyl$, $-(CH_2)_n-S-(CH_2)_m-R_7$, $-C(O)C(O)NH_2$, $-C(O)C(O)OR'_7$;

R_6 represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-alkynyl$, $-(CH_2)_m-S-(CH_2)_m-R_7$,


10



R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

15

R_{30} represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group, or




R_{32} and R_{61} , independently, represent small hydrophobic groups, preferably lower alkyls, and more preferably methyl;

Y_1 and Y_2 can, independently, be OH, or a group capable of being hydrolyzed to a hydroxyl group, including cyclic derivatives where Y_1 and Y_2 taken together form a ring having from 5 to 8 atoms in the ring structure,

m is zero or an integer in the range of 1 to 8; and

5 n is an integer in the range of 1 to 8.

22. The method of claim 1, wherein the inhibitor is represented by the general formula:




wherein:

10 R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl;

R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl;

R_3 represents hydrogen or an amino-protecting group;

15 R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or



20 R_6 represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-alkynyl$, $-(CH_2)_m-S-(CH_2)_m-R_7$;

R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;


U is absent or represents $-C(=O)-$, $-C(=S)-$, $-P(=O)(OR_8)-$, $-S(O_2)-$, or $-S(O)-$;

W represents a functional group which reacts with an active site residue of the targeted protease.

m is zero or an integer in the range of 1 to 8.

23. The method of claim 22, wherein:

W represents $-CN$, $-CH=NR_{53}$,



Y_1 and Y_2 are, independently, OH, or a group capable of being hydrolyzed to a hydroxyl group;

U represents $-C(=O)-$, $-C(=S)-$, or $-S(O_2)-$;

R_{50} represents O or S;

R_{51} represents N_3 , SH, NH_2 , NO_2 or OR'_7 ;

R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure;


R_{53} represents hydrogen, an alkyl, an alkenyl, an alkynyl, $-C(X_1)(X_2)-X_3$, $-(CH_2)_m-R_7$, $-(CH_2)_n-OH$, $-(CH_2)_n-O-alkyl$, $-(CH_2)_n-O-alkenyl$, $-(CH_2)_n-O-alkynyl$, $-(CH_2)_n-O-(CH_2)_m-R_7$, $-(CH_2)_n-SH$, $-(CH_2)_n-S-alkyl$, $-(CH_2)_n-S-alkenyl$, $-(CH_2)_n-S-alkynyl$, $-(CH_2)_n-S-(CH_2)_m-R_7$, $-C(O)C(O)NH_2$, $-C(O)C(O)OR'_7$;

X_1 represents a halogen;

X_2 and X_3 each represent a hydrogen or a halogen;


m is zero or an integer in the range of 1 to 8; and n is an integer in the range of 1 to 8.

24. The method of claim 1, wherein the inhibitor is represented by the general formula:



wherein:

W represents $-CN$, $-CH=NR_{53}$,



Y_1 and Y_2 are, independently, OH , or a group capable of being hydrolyzed to a hydroxyl group.

R_{50} represents O or S ;

R_{51} represents N_3 , SH , NH_2 , NO_2 or OR'_7 ;

R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the

phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure;

5 R_{53} represents hydrogen, an alkyl, an alkenyl, an alkynyl, $-C(X_1)(X_2)-X_3$, $-(CH_2)_m-R_7$, $-(CH_2)_n-OH$, $-(CH_2)_n-O-alkyl$, $-(CH_2)_n-O-alkenyl$, $-(CH_2)_n-O-alkynyl$, $-(CH_2)_n-O-(CH_2)_m-R_7$, $-(CH_2)_n-SH$, $-(CH_2)_n-S-alkyl$, $-(CH_2)_n-S-alkenyl$, $-(CH_2)_n-S-alkynyl$, $-(CH_2)_n-S-(CH_2)_m-R_7$, $-C(O)C(O)NH_2$, $-C(O)C(O)OR'_7$;

X_1 represents a halogen;

X_2 and X_3 each represent a hydrogen or a halogen;

10 m is zero or an integer in the range of 1 to 8; and


n is an integer in the range of 1 to 8.

25. The method of claim 24, wherein the Y_1 and Y_2 are connected via a ring having from 5 to 8 atoms in the ring structure.

15

26. The method of claim 25, wherein the Y_1 and Y_2 are pinacol.

27. The method of claim 24, wherein:



5 R_{33} represents hydrogen, an alkyl, an alkenyl, an alkynyl, $-C(X_1)(X_2)-X_3$, $-(CH_2)_m-R_7$, $-(CH_2)_n-OH$, $-(CH_2)_n-O-alkyl$, $-(CH_2)_n-O-alkenyl$, $-(CH_2)_n-O-alkynyl$, $-(CH_2)_n-O-(CH_2)_m-R_7$, $-(CH_2)_n-SH$, $-(CH_2)_n-S-alkyl$, $-(CH_2)_n-S-alkenyl$, $-(CH_2)_n-S-alkynyl$, $-(CH_2)_n-S-(CH_2)_m-R_7$, $-C(O)C(O)NH_2$, $-C(O)C(O)OR'_7$;

R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

10 R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

X_1 represents a halogen;

X_2 and X_3 , independently and for each occurrence, represent a hydrogen or a halogen;

15 m is zero or an integer in the range of 1 to 8; and n is an integer in the range of 1 to 8.


28. The method of claim 27, wherein:

X_1 represents fluorine;

X_2 and X_3 each independently represent a hydrogen or a fluorine.

20

29. The method of claim 24, wherein R_4 represents



wherein:

R₃₆ is a small hydrophobic group and R₃₈ is hydrogen, or, R₃₆ and R₃₈ together form a 4-7 membered heterocycle; and

5 R₄₀ represents a C-terminally linked amino acid residue or amino acid analog, or a C-terminally linked peptide or peptide analog, or an amino-protecting group.

30. The method of claim 24, wherein R₁ represents a lower alkyl.

10 31. The method of claim 30, wherein R₁ represents a methyl.

32. The method of claim 24, wherein R₂ represents a bulky hydrophobic group.

15 33. The method of claim 32, wherein the bulky hydrophobic group is a branched alkyl, branched alkenyl, branched alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, bicycloalkyl, bicycloalkenyl or bicycloalkynyl.

34. The method of claim 33, wherein the bulky hydrophobic group is a *t*-butyl or cyclohexyl.


20

35. The method of claim 24, wherein R₃ represents hydrogen.

36. The method of claim 24, wherein R₃₃ is a hydrogen, or a halogenated lower alkyl.

37. The method of claim 24, wherein X_1 is a fluorine, and X_2 and X_3 , if halogens, are fluorine.

5 38. The method of claim 24, represented by the general formula:




wherein;

10 R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl;

R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl;

R_3 represents hydrogen or an amino-protecting group;

15 R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or



20 R_6 represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O-alkyl$, $-(CH_2)_m-O-alkenyl$, $-(CH_2)_m-O-alkynyl$, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S-alkyl$, $-(CH_2)_m-S-alkenyl$, $-(CH_2)_m-S-alkynyl$, $-(CH_2)_m-S-(CH_2)_m-R_7$; R_7 represents, for each

occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

5 R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

10 R_{11} and R_{12} , independently each represent hydrogen, a alkyl, or a pharmaceutically acceptable salt, or R_{11} and R_{12} taken together with the O-B-O atoms to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure;

m is zero or an integer in the range of 1 to 8.


39. The method of claim 38, wherein:

15 R_1 represents methyl;

R_2 represents cyclohexyl;

R_3 , R_4 , R_{11} and R_{12} represent, independently and for each occurrence, hydrogen.

20 40. The method of claim 24, represented by the general formula:




wherein,

25 R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl;

R₂ represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl;

R₃ represents hydrogen or an amino-protecting group;

5 R₄ represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or




10 R_6 represents hydrogen, a halogen, an alkyl, an alkenyl, an alkynyl, an aryl, $-(CH_2)_m-R_7$, $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -alkyl, $-(CH_2)_m-O$ -alkenyl, $-(CH_2)_m-O$ -alkynyl, $-(CH_2)_m-O-(CH_2)_m-R_7$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -alkyl, $-(CH_2)_m-S$ -alkenyl, $-(CH_2)_m-S$ -alkynyl, $-(CH_2)_m-S-(CH_2)_m-R_7$; R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

15 R₇ represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R', represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle.

20 41. The method of claim 24, represented by the general formula:




wherein:

R_1 represents hydrogen, halogen or lower alkyl, lower alkenyl, or lower alkynyl;

R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl;

5 R_3 represents hydrogen or an amino-protecting group;

R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or




10 R_6 represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, $-(\text{CH}_2)_m\text{-}R_7$, $-(\text{CH}_2)_m\text{-OH}$, $-(\text{CH}_2)_m\text{-O-alkyl}$, $-(\text{CH}_2)_m\text{-O-alkenyl}$, $-(\text{CH}_2)_m\text{-O-alkynyl}$, $-(\text{CH}_2)_m\text{-O-}(\text{CH}_2)_m\text{-}R_7$, $-(\text{CH}_2)_m\text{-SH}$, $-(\text{CH}_2)_m\text{-S-alkyl}$, $-(\text{CH}_2)_m\text{-S-alkenyl}$, $-(\text{CH}_2)_m\text{-S-alkynyl}$, $-(\text{CH}_2)_m\text{-S-}(\text{CH}_2)_m\text{-}R_7$; R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

15 R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

20 X_1 , X_2 and X_3 each represent a hydrogen or a halogen.

42. The method of claim 24, wherein the (L)-Ala, (L)-Xaa-diastereomer of the inhibitor is represented by the general formula:




wherein:

R_2 represents a branched lower alkyl, aralkyl, aryl, heteroaralkyl, heteroaryl, cycloalkyl, or cycloalkylalkyl, e.g., optionally substituted by one or more substituents such as halogen, hydroxy, alkoxy, etc.,

R_3 represents hydrogen or an amino-protecting group;

R_4 represents hydrogen, a C-terminally linked amino acid residue or amino acid analog, a C-terminally linked peptide or peptide analog, an amino-protecting group, or




R_6 represents hydrogen, a halogen, a alkyl, a alkenyl, a alkynyl, an aryl, $-(\text{CH}_2)_m - \text{R}_7$, $-(\text{CH}_2)_m - \text{OH}$, $-(\text{CH}_2)_m - \text{O-alkyl}$, $-(\text{CH}_2)_m - \text{O-alkenyl}$, $-(\text{CH}_2)_m - \text{O-alkynyl}$, $-(\text{CH}_2)_m - \text{O} - (\text{CH}_2)_m - \text{R}_7$, $-(\text{CH}_2)_m - \text{SH}$, $-(\text{CH}_2)_m - \text{S-alkyl}$, $-(\text{CH}_2)_m - \text{S-alkenyl}$, $-(\text{CH}_2)_m - \text{S-alkynyl}$, $-(\text{CH}_2)_m - \text{S} - (\text{CH}_2)_m - \text{R}_7$;

R_7 represents, for each occurrence, a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

R'_7 represents, for each occurrence, hydrogen, or a substituted or unsubstituted alkyl, alkenyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle;

W represents $-\text{CN}$, $-\text{CH}=\text{NR}_{53}$.



Y_1 and Y_2 are, independently, OH, or a group capable of being hydrolyzed to a hydroxyl group;

5 R_{50} represents O or S;

R_{51} represents N_3 , SH, NH_2 , NO_2 or OR'_7 ;

R_{52} represents hydrogen, a lower alkyl, an amine, OR'_7 , or a pharmaceutically acceptable salt, or R_{51} and R_{52} taken together with the phosphorous atom to which they are attached complete a heterocyclic ring having from 5 to 8 atoms in the ring structure;

10 R_{53} represents hydrogen, an alkyl, an alkenyl, an alkynyl, $-\text{C}(\text{X}_1)(\text{X}_2)-\text{X}_3$, $-(\text{CH}_2)_m-\text{R}_7$, $-(\text{CH}_2)_n-\text{OH}$, $-(\text{CH}_2)_n-\text{O-alkyl}$, $-(\text{CH}_2)_n-\text{O-alkenyl}$, $-(\text{CH}_2)_n-\text{O-alkynyl}$, $-(\text{CH}_2)_n-\text{O}-(\text{CH}_2)_m-\text{R}_7$, $-(\text{CH}_2)_n-\text{SH}$, $-(\text{CH}_2)_n-\text{S-alkyl}$, $-(\text{CH}_2)_n-\text{S-alkenyl}$, $-(\text{CH}_2)_n-\text{S-alkynyl}$, $-(\text{CH}_2)_n-\text{S}-(\text{CH}_2)_m-\text{R}_7$, $-\text{C}(\text{O})\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{C}(\text{O})\text{OR}'_7$;

X_1 represents a halogen;

X_2 and X_3 each represent a hydrogen or a halogen;

m is zero or an integer in the range of 1 to 8; and

n is an integer in the range of 1 to 8.

20

43. The method of claim 42, wherein:



Y_1 and Y_2 represent independently each a hydroxyl;

R_2 represents cyclohexyl;

R_3 , R_4 , represent, independently and for each occurrence, hydrogen.





Figure 1

Figure 2**Synthesis Of Cyclohexylglycine boroAla**

DGS

4/19/01




Figure 3

Figure 4

MDS Project 003308




Figure 5

Percent Incidence of Diabetes





Figure 6



THIS PAGE BLANK (USPTO)